

Package ‘landsat8’

January 27, 2017

Type Package

Title Landsat 8 Imagery Rescaled to Reflectance, Radiance and/or Temperature

Version 0.1-10

Date 2017-01-26

Author Alexandre dos Santos

Maintainer Alexandre dos Santos <alexandre.santos@cas.ifmt.edu.br>

Description Functions for converted Landsat 8 multispectral satellite imagery rescaled to the top of atmosphere (TOA) reflectance, radiance and/or at satellite brightness temperature using radiometric rescaling coefficients provided in the metadata file (MTL file).

Imports rgdal, sp

Depends R (>= 3.1.2)

License GPL-2

LazyData TRUE

RoxygenNote 5.0.1

NeedsCompilation no

Repository CRAN

Date/Publication 2017-01-27 10:39:57

R topics documented:

| | |
|---------------------|---|
| band11 | 2 |
| band5 | 2 |
| radconv | 3 |
| reflconv | 4 |
| reflconvS | 5 |
| tempconv | 6 |

| | |
|--------------|----------|
| Index | 7 |
|--------------|----------|

band11

Sample Landsat 8 TIRS data

Description

GeoTIFF File Format containing a 300 x 300 pixel subset (1500 x 1500 m) of the Landsat 8 TIRS band 10 for path 228, row 71, obtained on 7 January 2011.

Usage

`data(band5)`

Format

GeoTIFF File Format format in 16 bits.

Source

Landsat images can be obtained from the United States Geological Survey at <http://landsat.usgs.gov>

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band11)
image(band11)
```

band5

Sample Landsat 8 OLI data

Description

GeoTIFF File Format containing a 300 x 300 pixel subset (1500 x 1500 m) of the Landsat 8 OLI band 5 for path 228, row 71, obtained on 7 January 2014.

Usage

`data(band5)`

Format

GeoTIFF File Format format in 16 bits.

Source

Landsat images can be obtained from the United States Geological Survey at <http://landsat.usgs.gov>

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band5)
image(band5)
```

radconv

Conversion to TOA Radiance

Description

Conversion to TOA radiance of satellite data.

Usage

```
radconv(x, M1, A1)
```

Arguments

| | |
|----|--|
| x | Image to be converted, in matrix, data frame, or SpatialGridDataFrame format. |
| M1 | band specific multiplicative rescaling factor from the metadata (MTL file) (RADIANCE_MULT_BAND_x, where x is the band number). |
| A1 | Mp band specific additive rescaling factor from the metadata (MTL file) (RADIANCE_ADD_BAND_x, where x is the band number). |

Value

TOA spectral radiance (Watts/(m2*srad*micro-m)).

Author(s)

Alexandre dos Santos

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band5)
band5.dn<- as(band5, 'SpatialGridDataFrame')
band5.rad<-radconv(band5.dn,5.9150E-03,-29.57525)
```

| | |
|----------|--------------------------------------|
| reflconv | <i>Conversion to TOA Reflectance</i> |
|----------|--------------------------------------|

Description

Conversion to TOA reflectance of satellite data.

Usage

```
reflconv(x, Mp, Ap)
```

Arguments

| | |
|----|---|
| x | Image to be converted, in matrix, data frame, or SpatialGridDataFrame format. |
| Mp | Band specific multiplicative rescaling factor from the metadata (MTL file) (REFLECTANCE_MULT_BAND_x, where x is the band number). |
| Ap | Band specific additive rescaling factor from the metadata (MTL file) (REFLECTANCE_ADD_BAND_x, where x is the band number). |

Value

TOA spectral radiance.

Author(s)

Alexandre dos Santos

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band5)
band5.dn<- as(band5, 'SpatialGridDataFrame')
band5.refl<-reflconv(band5.dn,2.0000E-05,-0.100000)
```

| | |
|-----------|--|
| reflconvS | <i>Conversion to TOA Reflectance with a Correction for the Sun Angle</i> |
|-----------|--|

Description

Conversion to TOA reflectance with a correction for the sun angle of satellite data.

Usage

```
reflconvS(x, Mp, Ap, sunelev)
```

Arguments

| | |
|---------|---|
| x | Image to be converted, in matrix, data frame, or SpatialGridDataFrame format. |
| Mp | band specific multiplicative rescaling factor from the metadata (MTL file) (REFLECTANCE_MULT_BAND_x, where x is the band number). |
| Ap | band specific additive rescaling factor from the metadata (MTL file) (REFLECTANCE_ADD_BAND_x, where x is the band number). |
| sunelev | Sun elevation in degrees is provided in the metadata (MTL file) (SUN_ELEVATION). |

Value

TOA spectral radiance with a correction for the sun angle.

Author(s)

Alexandre dos Santos

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band5)
band5.dn<- as(band5, 'SpatialGridDataFrame')
band5.reflS<-reflconvS(band5.dn,2.0000E-05,-0.100000,41.12846745)
```

`tempconv`*Conversion to At Satellite Brightness Temperature*

Description

Conversion to At satellite brightness temperature of satellite data.

Usage

```
tempconv(x, M1, A1, K1, K2)
```

Arguments

| | |
|-----------------|---|
| <code>x</code> | Image to be converted, in matrix, data frame, or <code>SpatialGridDataFrame</code> format. |
| <code>M1</code> | band specific multiplicative rescaling factor from the metadata (MTL file) (<code>RADIANCE_MULT_BAND_x</code> , where <code>x</code> is the band number). |
| <code>A1</code> | <code>Mp</code> band specific additive rescaling factor from the metadata (MTL file) (<code>RADIANCE_ADD_BAND_x</code> , where <code>x</code> is the band number). |
| <code>K1</code> | band specific thermal conversion constant from the metadata (MTL file) (<code>K1_CONSTANT_BAND_x</code> , where <code>x</code> is the band number, 10 or 11). |
| <code>K2</code> | band specific thermal conversion constant from the metadata (MTL file) (<code>K2_CONSTANT_BAND_x</code> , where <code>x</code> is the band number, 10 or 11). |

Value

At satellite brightness temperature in Kelvin (K).

Author(s)

Alexandre dos Santos

References

U.S. Geological Survey. 2015. Landsat 8 (L8) data users handbook. Version 1.0. 97p.

Examples

```
data(band11)
band11.dn<- as(band11, 'SpatialGridDataFrame')
band11.tempK<-tempconv(band11.dn, 3.3420E-04, 0.10000, 480.89, 1201.14)
```

Index

*Topic **datasets**

band11, [2](#)

band5, [2](#)

band11, [2](#)

band5, [2](#)

radconv, [3](#)

reflconv, [4](#)

reflconvS, [5](#)

tempconv, [6](#)