

Package ‘seacarb’

October 18, 2021

Title Seawater Carbonate Chemistry

Version 3.3.0

Date 2021-10-15

LazyData true

Depends R (>= 3.5.0), oce, gsw, SolveSAPHE

Description Calculates parameters of the seawater carbonate system and assists the design of ocean acidification perturbation experiments.

Encoding UTF-8

URL <https://CRAN.R-project.org/package=seacarb>

License GPL (>= 2)

Repository CRAN

RoxygenNote 7.1.1

NeedsCompilation no

Author Jean-Pierre Gattuso [aut, cre, cph],
Jean-Marie Epitalon [aut],
Heloise Lavigne [aut],
James Orr [aut],
Bernard Gentili [ctb],
Mathilde Hagens [ctb],
Andreas Hofmann [ctb],
Jens-Daniel Mueller [ctb],
Aurélien Proye [ctb],
James Rae [ctb],
Karline Soetaert [ctb]

Maintainer Jean-Pierre Gattuso <jean-pierre.gattuso@imev-mer.fr>

Date/Publication 2021-10-18 11:50:02 UTC

R topics documented:

alkalinity	3
amp	4

bjerrum	5
bor	7
buffer	8
buffergen	13
buffesm	18
carb	23
carbb	29
carbfull	35
d2p	41
derivnum	42
eos2teos_chem	47
eos2teos_geo	48
errors	50
f2pCO2	57
fCO2insi	58
K0	59
K1	60
K1p	63
K2	64
K2p	67
K2si	68
K3p	69
Kb	71
kconv	72
Kf	74
kfg	75
Khs	76
Kn	77
Ks	78
Ksi	79
Kspa	80
Kspc	82
Kw	83
oa	84
Om	89
p2d	93
p2fCO2	94
p2xCO2	95
pCa	96
pCO2insi	101
Pcoeffs	102
Pcorrect	103
pgas	104
pH	109
pHconv	110
pHinsi	111
pHslope	115
pHspec	116

pmix	117
ppH	121
psi	126
pTA	130
rho	135
sa2sp_chem	135
sa2sp_geo	137
seacarb_test_P0	138
seacarb_test_P300	139
seaFET	139
sf_calc	140
sf_calib	141
sir	143
sir_b	149
sir_full	155
sp2sa_chem	161
sp2sa_geo	163
speciation	164
teos2eos_chem	165
teos2eos_geo	166
theta	168
tris	169
vapress	170
x2pCO2	171

alkalinity	<i>Example data file for function at</i>
------------	--

Description

The variables are:

- volume: Volume of acid added to the sample in ml
- E: Potential measured during the titration in mV
- temperature: Temperature in degrees Celsius
- weight: Weight of the sample in g
- S: Salinity
- normality : Normality of the acid
- ETris: Potential used for the calibration of the electrode in mV
- pHTris: pH used for the calibration of the electrode with the TRIS buffer

Usage

alkalinity

Format

A data frame with 29 rows and 8 variables

Source

Data come from a potentiometric titration performed by Steeve Comeau.

amp	<i>pH value of the AMP buffer</i>
-----	-----------------------------------

Description

pH value of the AMP buffer (on the total scale in mol/kg)

Usage

`amp(S=35, T=25)`

Arguments

S Salinity, default is 35

T Temperature in degrees Celsius, default is 25oC

Details

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

AMP pH value of the AMP buffer (on the total scale in mol/kg)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication 3*, 1-191.

See Also

[tris](#), [pHslope](#), [pH](#).

Examples

```
##Example from Dickson et al. (2007)
amp(S=35,T=25)
```

bjerrum

Bjerrum plot

Description

Plot the concentration of the various ionic forms of a molecule as a function of pH

Usage

```
bjerrum(K1=K1(), K2=NULL, K3=NULL, phmin=2, phmax=12, by=0.1, conc=1,
        type="l", col="black", ylab="Relative concentration (%)", add=FALSE, ...)
```

Arguments

K1	First dissociation constant
K2	Second dissociation constant, default is NULL
K3	Third dissociation constant, default is NULL
phmin	Minimum pH value, default is 2
phmax	Maximum pH value, default is 12
by	Increment on the pH axis, default is 0.1
conc	concentration of molecule, default is 1
type	Type of plot, default is line
col	Color of plot, default is black
ylab	Label of Y axis, default is (mol/kg)
add	false:start new, true: add to current, default is false
...	Graphical parameters (see par) and any further arguments of plot, typically plot.default , may also be supplied as arguments to this function. Hence, the high-level graphics control arguments described under par and the arguments to title may be supplied to this function.

Details

Note that the concentration is plotted in mol/kg only if conc is given in mol/kg

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl>

References

Zeebe, R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

See Also

[matplot](#), [par](#), [speciation](#).

Examples

```
## Plot the bjerrum plot for the carbonate system using the default values
bjerrum(K1(),K2(),main="DIC speciation",lwd=2)
abline(v=-log10(K1()),col="grey")
mtext(side=3,at=-log10(K1()),"pK1")
abline(v=-log10(K2()),col="grey")
mtext(side=3,at=-log10(K2()),"pK2")
legend("left",lty=1:3,lwd=2,legend=c(expression(CO[2]),expression(HCO[3]^"-"),
expression(CO[3]^"2-")))

## Plot the bjerrum plot for phosphate using the default values
bjerrum(K1p(),K2p(),K3p(),main="phosphate speciation",lwd=2)
legend("left",lty=1:4,lwd=2,legend=c(expression(H[3]~PO[4]),
expression(H[2]~PO[4]^"-"),
expression(HPO[4]^"2-"),expression(PO[4]^"3-")))

## Plot the bjerrum plot for the carbonate system using the values other
## than the default ones, showing the effect of temperature
bjerrum(K1(T=25,S=35),K2(T=25,S=35),conc=1.3,main="effect of temperature" )
bjerrum(K1(T=0,S=35),K2(T=0,S=35),conc=1.3,add=TRUE,col="red")
legend("left",lty=1,col=c("black","red"),legend=c("T=25 oC","T=0 oC"))
legend("right",lty=1:3,legend=c(expression(CO[2]),expression(HCO[3]^"-"),
expression(CO[3]^"2-")))

## Plot the bjerrum plot for the carbonate system using the values other
## than the default ones, showing the effect of salinity
bjerrum(K1(T=25,S=35),K2(T=25,S=35),conc=1.3,main="effect of salinity" )
bjerrum(K1(T=25,S=5),K2(T=25,S=5),conc=1.3,add=TRUE,col="blue")
legend("left",lty=1,col=c("black","blue"),legend=c("S=35","S=5"))
legend("right",lty=1:3,legend=c(expression(CO[2]),expression(HCO[3]^"-"),
expression(CO[3]^"2-")))

## Plot the bjerrum plot for the carbonate system using the values other
## than the default ones, showing the effect of pressure
bjerrum(K1(P=0),K2(P=0),conc=1.3,main="effect of pressure" )
bjerrum(K1(P=300),K2(P=300),conc=1.3,add=TRUE,col="green")
legend("left",lty=1,col=c("black","green"),legend=c("P=0","P=300"),title="atm")
legend("right",lty=1:3,legend=c(expression(CO[2]),expression(HCO[3]^"-"),
expression(CO[3]^"2-")))
```

bor	<i>Total boron concentration (mol/kg)</i>
-----	---

Description

total boron concentration ($mol\ kg^{-1}$)

Usage

bor(S, b)

Arguments

S	Salinity, default is 35
b	"l10" for using the formulation of Lee et al. (2010), "u74" for using the Uppstrom (1974), or "k18" for using the Kulinski et al. (2018), default is "u74"

Details

Note that the formulation of Kulinski et al. (2018) is specifically designed for the Baltic Sea. Three formulations are described in their paper:

- based on their measurements: $TB = [\mu\text{mol/kg}] = 10.838 * S + 13.821$
- based on Kremling (1970 and 1972): $TB [\mu\text{mol/kg}] = 11.44 * S + 12.6$; $R^2 = 0.95$
- consensus regression (Kremling + their data): $TB [\mu\text{mol/kg}] = 11.405 * S + 11.869$; $R^2 = 0.97s$

The latter formulation is used here.

Value

bor total boron concentration ($mol\ kg^{-1}$)

Author(s)

Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Kulinski K., Szymczyska B., Koziorowska K., Hammer K. & Schneider B., 2018. Anomaly of total boron concentration in the brackish waters of the Baltic Sea and its consequence for the CO₂ system calculations. *Marine Chemistry*. doi:s10.1016/j.marchem.2018.05.007.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantic oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Examples

```
bor(35, "110")
```

buffer

Buffer parameters of the seawater carbonate system

Description

Returns buffer parameters of the seawater carbonate system.

Usage

```
buffer(flag, var1, var2, S = 35, T = 25, Patm = 1, P = 0, Pt = 0, Sit = 0,
      k1k2 = "x", kf = "x", ks = "d", pHscale = "T", b = "u74", warn = "y",
      eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
------	--

var1	enter value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	enter value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.

- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

PhiD	PhiD, chemical buffer factor ($d\text{pH}/d[\text{DIC}]$); input/output of dissolved CO ₂ (unit pH per mol/kg)
BetaD	BetaD, homogeneous or Revelle buffer factor ($d\ln(\text{pCO}_2)/d\ln[\text{DIC}]$); input/output of dissolved CO ₂
PiD	PiD, chemical buffer factor ($d\text{pCO}_2/d[\text{DIC}]$); input/output of dissolved CO ₂ (μatm per mol/kg)
PhiB	PhiB, chemical buffer factor ($d\text{pH}/d[\text{DIC}]$); from input/output of bicarbonate (unit pH per mol/kg)
BetaB	BetaB, homogeneous buffer factor ($d\ln(\text{pCO}_2)/d\ln[\text{DIC}]$); input/output of bicarbonate
PiB	PiB, chemical buffer factor ($d\text{pCO}_2/d[\text{DIC}]$); input/output of dissolved CO ₂ (μatm per mol/kg)
PhiC	PhiC, chemical buffer factor ($d\text{pH}/d[\text{DIC}]$); input/output of carbonate (unit pH per mol/kg)
BetaC	BetaC, homogeneous buffer factor ($d\ln(\text{pCO}_2)/d\ln[\text{DIC}]$); input/output of carbonate
PiC	PiC, chemical buffer factor ($d\text{pCO}_2/d[\text{DIC}]$); input/output of carbonate (μatm per mol/kg)
PhiH	PhiH, chemical buffer factor ($d\text{pH}/d[\text{ALK}]$); input/output of strong acid (unit pH per mol/kg)
PiH	PiH, chemical buffer factor ($d\text{pCO}_2/d[\text{ALK}]$); input/output of strong acid (μatm per mol/kg)

Author(s)

Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Frankignoulle M., 1994 A complete set of buffer factors for acid/base CO_2 system in seawater. *Journal of Marine Systems* **5**, 111-118.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Examples

```
## Computation with a couple of variables
buffer(flag=8, var1=8.2, var2=0.00234, S=35, T=25, Patm=1, P=0, Pt=0,
      S1t=0, pHscale="T", kf="pf", k1k2="1", b="u74")

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
```

```

Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("1", "1", "1")
pHscale <- c("T", "T", "T")
b <- c("110", "110", "110")
buffer(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt,
Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

## Test for all flags

flag <- c(1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 21, 22, 23, 24, 25)

var1 <- c(8.200000, 7.477544e-06, 7.477544e-06, 7.477544e-06, 7.477544e-06, 8.2,
8.2, 8.2, 8.2, 0.001685024, 0.001685024, 0.001685024, 0.0002888382,
0.0002888382, 0.002391252, 264.2008, 264.2008, 264.2008, 264.2008, 264.2008)

var2 <- c(7.477544e-06, 0.001685024, 0.0002888382, 0.002391252, 0.001981340,
0.001685024, 0.0002888382, 0.002391252, 0.001981340, 0.0002888382, 0.002391252,
0.001981340, 0.002391252, 0.001981340, 0.001981340, 8.2, 0.001685024,
0.0002888382, 0.002391252, 0.001981340)

buffer(flag=flag, var1=var1, var2=var2)

```

buffergen

Buffer factors of the seawater carbonate system as defined by Hagens and Middelburg (2016)

Description

Returns the suite of buffer factors presented in Table 3 of Hagens and Middelburg (2016), as well as the proton concentration buffer factor (beta.H of Hofmann et al, 2010) and the classic Revelle factor. For practical purposes, this function excludes the nitrate and nitrite acid-base systems presented in this paper, as well as the fully protonated form of sulfate (H₂SO₄) and fully deprotonated form of sulfide (S²⁻), as their contributions to total alkalinity under natural seawater conditions are negligible. Its input arguments are identical to those in the carbfull function of seacarb.

Usage

```
buffergen(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0, k1k2="x", kf="x",
ks="d", pHscale="T", b="u74", gas="potential", NH4t=0, HSt=0)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given
------	--

	flag = 3 CO2 and CO3 given
	flag = 4 CO2 and ALK given
	flag = 5 CO2 and DIC given
	flag = 6 pH and HCO3 given
	flag = 7 pH and CO3 given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO3 and CO3 given
	flag = 11 HCO3 and ALK given
	flag = 12 HCO3 and DIC given
	flag = 13 CO3 and ALK given
	flag = 14 CO3 and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "mp2" from Mojica Prieto et al. (2002), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "w14" from Waters et al. (2014), "p18" from Papadimitriou et al. (2018), "s20" from Sulpis et al. (2020), "sb21" from Shockman & Byrne (2021), and "r" from Roy et al. (1993). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)

b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO ₂ , i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO ₂ is not an input variable (flags 1 to 15). The default is "potential" and should be a unique value..
NH4t	Concentration of total ammonium in mol/kg; set to 0 if NA
HSt	Concentration of total hydrogen sulfide in mol/kg; set to 0 if NA

Details

The Lueker et al. (2000) constants for K₁ and K₂, the Perez and Fraga (1987) constant for K_f and the Dickson (1990) constant for K_s are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K₁ and K₂:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K₁ and K₂ formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K₁ and K₂ are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K₂, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K₁, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K₁ and K₂ formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K₁ and K₂ are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For K_f:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For K_s:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.

- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs, Ks_i and K2s_i, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_f, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n, the pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

The function returns a list containing the following matrices:

Carbfull	Output of the carbfull function that is used within buffergen
dALK.dH	Sensitivity of ALK to a change in proton concentration (dimensionless). Species-specific.
dtotX.dH	Sensitivity of an acid-base species to a change in proton concentration (dimensionless). Species-specific.
dALK.dX	Sensitivity of ALK to a change in an acid-base species (dimensionless). Species-specific.
dtotX.dX	Sensitivity of an acid-base species to a change in its total concentration (dimensionless). Species-specific.
dALK.dpH	Sensitivity of ALK to a change in pH (mol/kg-soln). Species-specific.
dtotX.dpH	Sensitivity of an acid-species to a change in pH (mol/kg-soln). Species-specific.
dH.dALK	Sensitivity of proton concentration to a change in ALK (dimensionless). Values are the same for all species and all acid-base systems, except for the fluoride and sulfate acid-base systems, which slightly deviate due to pH scale conversion effects.

dH.dtotX	Sensitivity of an acid-species to a change in its total concentration (dimensionless). Values are the same for all species of a specific acid-base system.
dX.dALK	Sensitivity of an acid-species to a change in its total concentration (dimensionless). Species-specific.
dX.dtotX	Sensitivity of an acid-species to a change in its total concentration (dimensionless). Species-specific.
dpH.dALK	Sensitivity of pH due to a change in ALK ((mol/kg-soln)-1). Values are the same for all species and all acid-base systems, except for the fluoride and sulfate acid-base systems, which slightly deviate due to pH scale conversion effects.
dpH.dtotX	Sensitivity of pH due to a change in the total concentration of an acid-base system ((mol/kg-soln)-1). Values are the same for all species of a specific acid-base system.
beta.H	proton concentration buffer factor (Eq.4 of Hagens and Middelburg (2016), dimensionless)
RF	Revelle factor (dimensionless)

If the total concentration of an acid-base system is 0, the values of the buffer factors corresponding to all species of that acid-base system return NA.

Author(s)

Mathilde Hagens <m.hagens@uu.nl>

References

Hagens M. and Middelburg J. J., 2016 Generalised expressions for the response of pH to changes in ocean chemistry. *Geochimica et Cosmochimica Acta* **187** 334-349.

Examples

```
## With a couple of variables
buffergen(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="1", ks="d", b="u74", gas="potential", NH4t=0, HSt=0)

## With a couple of variables and non-zero nutrient concentrations
buffergen(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=5e-6, Sit=2e-6,
pHscale="T", kf="pf", k1k2="1", ks="d", b="u74", gas="potential", NH4t=10e-6, HSt=0.1e-6)

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
```

```

k1k2 <- c("1", "1", "1")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
gas <- c("potential")
NH4t <- c(0, 0, 0)
HSt <- c(0, 0, 0)
buffergen(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt, Sit=Sit,
kf=kf, k1k2=k1k2, pHscale=pHscale, b=b, gas=gas, NH4t=NH4t, HSt=HSt)

## Test with all flags
flag <- c((1:15), (21:25))
var1 <- c(8.200000, 7.308171e-06, 7.308171e-06, 7.308171e-06, 7.308171e-06,
8.2, 8.2, 8.2, 8.2, 0.001646857, 0.001646857, 0.001646857, 0.0002822957,
0.0002822957, 0.00234, 258.2164, 258.2164, 258.2164, 258.2164, 258.2164 )
var2 <- c(7.308171e-06, 0.001646857, 0.0002822957, 0.00234, 0.001936461,
0.001646857, 0.0002822957, 0.00234, 0.001936461, 0.0002822957,
0.00234, 0.001936461, 0.00234, 0.001936461, 0.001936461, 8.2,
0.001646857, 0.0002822957, 0.00234, 0.001936461)
buffergen(flag=flag, var1=var1, var2=var2)

```

buffesm

Buffer capacities of the seawater carbonate system from Egleston et al. (2010), corrected and enhanced

Description

Returns the six buffer factors of the seawater carbonate system as defined by Egleston, Sabine and Morel (2010), denoted here as ESM. Also returns the classic Revelle factor (relative change in pCO₂ over that for DIC). In ESM, there are errors in the equations in Table 1 for S , Ω_{DIC} , and Ω_{Alk} . These errors have been corrected here. The results of this routine have been validated: when input concentrations of Pt and Sit are set to zero, they produce results that are identical to those shown in ESM's Fig. 2. But when Pt and Sit are nonzero, contributions from phosphoric and silicic acid systems are taken into account, an improvement to the Egleston et al. (2010) approach. This routine was inspired and adapted from seacarb's "buffer" function. Its input arguments are identical to those in the "buffer" and "carb" functions of seacarb.

Usage

```
buffesm(flag, var1, var2, S = 35, T = 25, Patm = 1, P = 0, Pt = 0, Sit = 0,
k1k2 = "x", kf = "x", ks = "d", pHscale = "T", b = "u74", warn = "y",
eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given
------	--

	flag = 4 CO2 and ALK given
	flag = 5 CO2 and DIC given
	flag = 6 pH and HCO3 given
	flag = 7 pH and CO3 given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO3 and CO3 given
	flag = 11 HCO3 and ALK given
	flag = 12 HCO3 and DIC given
	flag = 13 CO3 and ALK given
	flag = 14 CO3 and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	enter value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	enter value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA; when nonzero, account for phosphoric acid system, unlike in Egleston et al. (2010).
Sit	Concentration of total silicate in mol/kg; set to 0 if NA; when nonzero, account for silicic acid system, unlike in Egleston et al. (2010).
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)

b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For K0:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

gammaDIC	γ_{DIC} , ocean's capacity to buffer changes in [CO ₂] due to accumulation of CO ₂ from the atmosphere $(\partial \ln[CO_2]/\partial DIC)^{-1}$ (units = mol/kg; multiply by 1000 to get mmol/kg, i.e., the units presented in Egleston et al., 2010)
betaDIC	β_{DIC} , ocean's capacity to buffer changes in [H ⁺] due to accumulation of CO ₂ from the atmosphere $(\partial \ln[H^+]/\partial DIC)^{-1}$ (units = mol/kg)
omegaDIC	Ω_{DIC} , ocean's capacity to buffer changes in [CO ₃ ²⁻] due to accumulation of CO ₂ from the atmosphere $(\partial \ln[CO_3^{2-}]/\partial DIC)^{-1}$; same as $(\partial \ln\Omega_A/\partial DIC)^{-1}$ and $(\partial \ln\Omega_C/\partial DIC)^{-1}$ (units= mol/kg)
gammaALK	γ_{ALK} , ocean's capacity to buffer changes in [CO ₂] due to changes in alkalinity $(\partial \ln[CO_2]/\partial ALK)^{-1}$ (units = mol/kg)
betaALK	β_{ALK} , ocean's capacity to buffer changes in [H ⁺] due to changes in alkalinity $(\partial \ln[H^+]/\partial ALK)^{-1}$ (units = mol/kg)

omegaALK	Ω_{ALK} , ocean's capacity to buffer changes in $[CO_3^{2-}]$ due to changes in alkalinity $(\partial \ln [CO_3^{2-}] / \partial ALK)^{-1}$; same as $(\partial \ln \Omega_A / \partial ALK)^{-1}$ and $(\partial \ln \Omega_C / \partial ALK)^{-1}$ (units = mol/kg)
R	Revelle factor, relative change in $[CO_2]$ or pCO_2 over the relative change in DIC $(\partial \ln [CO_2] / \partial \ln DIC)^{-1}$ (unitless)

Author(s)

James Orr <James.Orr@lsce.ipsl.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $AgCl(s) + 1/2H_2(g) = Ag(s) + HCl(aq)$, and the standard acidity constant of the ion HSO_4 in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Egleston, E. S., Sabine, C. L. and Morel, F. M. M., 2010 Revelle revisited: Buffer factors that quantify the response of ocean chemistry to changes in DIC and alkalinity, *Global Biogeochem. Cycles* **24**, GB1002, doi:10.1029/2008GB003407.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Frankignoulle M., 1994 A complete set of buffer factors for acid/base CO_2 system in seawater. *Journal of Marine Systems* **5**, 111-118.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Examples

```

## Computation with a couple of variables
buffesm(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Pt=0,
Sit=0, pHscale="T", kf="pf", k1k2="l", b="u74")

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("l", "l", "l")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
buffesm(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt,
Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

## Test for all flags
flag <- c(1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 21, 22, 23, 24, 25)

var1 <- c(8.200000, 7.477544e-06, 7.477544e-06, 7.477544e-06, 7.477544e-06, 8.2,
8.2, 8.2, 8.2, 0.001685024, 0.001685024, 0.001685024, 0.0002888382,
0.0002888382, 0.002391252, 264.2008, 264.2008, 264.2008, 264.2008, 264.2008)
var2 <- c(7.477544e-06, 0.001685024, 0.0002888382, 0.002391252, 0.001981340,
0.001685024, 0.0002888382, 0.002391252, 0.001981340, 0.0002888382, 0.002391252,
0.001981340, 0.002391252, 0.001981340, 0.001981340, 8.2, 0.001685024,
0.0002888382, 0.002391252, 0.001981340)
buffesm(flag=flag, var1=var1, var2=var2)

## Compute 2 additional factors of interest (ratios of relative changes)
be <- buffesm(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt,
Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)
#      Ratio of gammaDIC/betaDIC = d ln [H+] / d ln pCO2
Hfac <- (be$gammaDIC/be$betaDIC)                                #H+ factor
#      Ratio of gammaDIC/omegaDIC = d ln [CO32-] / d ln pCO2
Satfac <- (be$gammaDIC/be$omegaDIC)                                #Saturation factor

```

Description

Returns parameters of the seawater carbonate system.

Usage

```
carb(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
      k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential",
      warn="y", eos="eos80", long=1.e20, lat=1.e20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".

kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.

- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μ atm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μ atm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μ atm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μ atm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
fCO2insitu	"in situ" fCO2, CO2 fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
HC03	HC03 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Heloise Lavigne, James Orr and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* 7, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21**, 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO_2 in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```

## With a couple of variables
carb(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74")

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("l", "l", "l")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
carb(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P,
Pt=Pt, Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

## Test with all flags
flag <- c((1:15), (21:25))
var1 <- c(8.200000, 7.308171e-06, 7.308171e-06, 7.308171e-06, 7.308171e-06,
8.2, 8.2, 8.2, 8.2, 0.001646857, 0.001646857, 0.001646857, 0.0002822957,
0.0002822957, 0.00234, 258.2164, 258.2164, 258.2164, 258.2164, 258.2164 )
var2 <- c(7.308171e-06, 0.001646857, 0.0002822957, 0.00234, 0.001936461,
0.001646857, 0.0002822957, 0.00234, 0.001936461, 0.0002822957,
0.00234, 0.001936461, 0.00234, 0.001936461, 0.001936461, 8.2,
0.001646857, 0.0002822957, 0.00234, 0.001936461)
carb(flag=flag, var1=var1, var2=var2)

## Test using a data frame
data(seacarb_test_P0) #test data set for P=0 (surface)
tab <- seacarb_test_P0[14:19,]

## method 1 using the column numbers
carb(flag=tab[[1]], var1=tab[[2]], var2=tab[[3]], S=tab[[4]], T=tab[[5]],
P=tab[[6]], Sit=tab[[8]], Pt=tab[[7]])

## method 2 using the column names
carb(flag=tab$flag, var1=tab$var1, var2=tab$var2, S=tab$S, T=tab$T,
P=tab$P, Sit=tab$Sit, Pt=tab$Pt)

```

Description

Returns parameters of the seawater carbonate system when boron is added.

Usage

```
carbb(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
      k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential", badd=0,
      warn="y", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA

k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
badd	Amount of boron added in mol/kg.
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.

- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μ atm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μ atm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μ atm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μ atm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
fCO2insitu	"in situ" fCO2, CO2 fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
HCO3	HCO3 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Heloise Lavigne, James Orr and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl}(\text{s}) + 1/2\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34. Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```
## With a couple of variables
carbb(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74", badd=0)
```

carbfull

Parameters of the seawater carbonate system - extension of carb

Description

Returns parameters of the seawater carbonate system, including the ammonium and sulfide acid-base systems, as well as full acid-base speciation

Usage

```
carbfull(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential",
NH4t=0, HSt=0)
```

Arguments

flag	select the couple of variables available. The flags which can be used are:
	flag = 1 pH and CO ₂ given
	flag = 2 CO ₂ and HCO ₃ given
	flag = 3 CO ₂ and CO ₃ given
	flag = 4 CO ₂ and ALK given
	flag = 5 CO ₂ and DIC given
	flag = 6 pH and HCO ₃ given
	flag = 7 pH and CO ₃ given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO ₃ and CO ₃ given
	flag = 11 HCO ₃ and ALK given
	flag = 12 HCO ₃ and DIC given
	flag = 13 CO ₃ and ALK given
	flag = 14 CO ₃ and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO ₂ and pH given
	flag = 22 pCO ₂ and HCO ₃ given

	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg-soln, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg-soln, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg-soln; set to 0 if NA
Sit	Concentration of total silicate in mol/kg-soln; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "mp2" from Mojica Prieto et al. (2002), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "w14" from Waters et al. (2014), "p18" from Papadimitriou et al. (2018), "s20" from Sulpis et al. (2020), "sb21" from Shockman & Byrne (2021), and "r" from Roy et al. (1993). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
NH4t	Concentration of total ammonium in mol/kg-soln; set to 0 if NA
HSt	Concentration of total hydrogen sulfide in mol/kg-soln; set to 0 if NA

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs, Ks and K2si, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , the pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg-soln)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μ atm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μ atm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μ atm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μ atm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
fCO2insitu	"in situ" fCO2, CO2 fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
HCO3	HCO3 concentration (mol/kg-soln)
CO3	CO3 concentration (mol/kg-soln)
DIC	DIC concentration (mol/kg-soln)
ALK	ALK, total alkalinity (mol/kg-soln)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state
NH4	NH4 concentration (mol/kg-soln)
NH3	NH3 concentration (mol/kg-soln)
BOH3	B(OH)3 concentration (mol/kg-soln)
BOH4	B(OH)4 concentration (mol/kg-soln)
H3PO4	H3PO4 concentration (mol/kg-soln)
H2PO4	H2PO4 concentration (mol/kg-soln)

HP04	HPO4 concentration (mol/kg-soln)
P04	PO4 concentration (mol/kg-soln)
H2S	H2S concentration (mol/kg-soln)
HS	HS concentration (mol/kg-soln)
SiOH4	SiOH4 concentration (mol/kg-soln)
SiOOH3	SiOOH3 concentration (mol/kg-soln)
SiO2OH2	SiO2OH2 concentration (mol/kg-soln)
HF	HF concentration (mol/kg-soln)
F	F concentration (mol/kg-soln)
HSO4	HSO4 concentration (mol/kg-soln)
SO4	SO4 concentration (mol/kg-soln)
H	H concentration at specified pH scale (mol/kg-soln)
OH	OH concentration (mol/kg-soln)
NH4t	Supplied NHt concentration (mol/kg-soln); values are recycled if necessary
B0R	Calculated total borate concentration (mol/kg-soln)
Pt	Supplied Pt concentration (mol/kg-soln); values are recycled if necessary
HSt	Supplied HSt concentration (mol/kg-soln); values are recycled if necessary
Sit	Supplied Sit concentration (mol/kg-soln); values are recycled if necessary
FLUO	Calculated total fluoride concentration (mol/kg-soln)
ST	Calculated total sulfate concentration (mol/kg-soln)

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2014)

Author(s)

Heloise Lavigne, James Orr, Mathilde Hagens and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **22**, vol49 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2014. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences Discussions* **11**, 5327-5397.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], Mar. Chem. 165, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Mar. Chem.*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```
## With a couple of variables
carbfull(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74", gas="potential", NH4t=0, HSt=0)

## With a couple of variables and non-zero nutrient concentrations
carbfull(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=5e-6, Sit=2e-6,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74", gas="potential", NH4t=10e-6, HSt=0.1e-6)

## Using vectors as arguments
```

```

flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("l", "l", "l")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
gas <- c("potential", "potential", "potential")
NH4t <- c(0, 0, 0)
HSt <- c(0, 0, 0)
carbfull(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt, Sit=Sit,
kf=kf, k1k2=k1k2, pHscale=pHscale, b=b, gas=gas, NH4t=NH4t, HSt=HSt)

## Test with all flags
flag <- c((1:15), (21:25))
var1 <- c(8.200000, 7.308171e-06, 7.308171e-06, 7.308171e-06, 7.308171e-06,
8.2, 8.2, 8.2, 8.2, 0.001646857, 0.001646857, 0.001646857, 0.0002822957,
0.0002822957, 0.00234, 258.2164, 258.2164, 258.2164, 258.2164, 258.2164 )
var2 <- c(7.308171e-06, 0.001646857, 0.0002822957, 0.00234, 0.001936461,
0.001646857, 0.0002822957, 0.00234, 0.001936461, 0.0002822957,
0.00234, 0.001936461, 0.00234, 0.001936461, 0.001936461, 8.2,
0.001646857, 0.0002822957, 0.00234, 0.001936461)
carbfull(flag=flag, var1=var1, var2=var2)

```

d2p

*Converts depth in meters to pressure in dbar***Description**

Converts depth in meters to pressure in dbar

Usage

```
d2p(depth, lat=40)
```

Arguments

depth	Depth in meters
lat	Latitude in degrees, N and S is irrelevant, default is 40o

Value

pressure	Pressure corresponding to the depth given, in dbar
----------	--

Author(s)

Heloise Lavigne and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Saunders P. M., 1981. Practical conversion of pressure to depth. *J. Phys. Oceanogr.* **11**: 573-574.

See Also

[p2d](#)

Examples

```
d2p(depth=7500, lat=30)
```

derivnum

Numerical derivatives of seawater carbonate system variables

Description

Returns numerical derivatives of the seawater carbonate system output variables with respect to input variables.

Usage

```
derivnum(varid, flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential", warn="y",
eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

varid	Variable length, case insensitive, character identifier of variable with respect to which derivatives are requested. Possible values are: '1' or 'var1' : Variable 1 of the input pair (This is TAlk if flag is 15) '2' or 'var2' : Variable 2 of the input pair (This is DIC if flag is 15) 'sil', 'silt', 'tsil' or 'silicate' : Total silicate concentration 'phos', 'phost', 'tphos' or 'phosphate' : Total phosphate concentration 't', 'temp' or 'temperature' : temperature 's', 'sal' or 'salinity' : salinity 'K0','K1','K2','Kb','Kw','Kspa' or 'Kspc' : one of the dissociation constants 'bor' : total boron
flag	select the input pair of carbonate-system variables available by choosing the following flag: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given

	flag = 3 CO ₂ and CO ₃ given
	flag = 4 CO ₂ and ALK given
	flag = 5 CO ₂ and DIC given
	flag = 6 pH and HCO ₃ given
	flag = 7 pH and CO ₃ given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO ₃ and CO ₃ given
	flag = 11 HCO ₃ and ALK given
	flag = 12 HCO ₃ and DIC given
	flag = 13 CO ₃ and ALK given
	flag = 14 CO ₃ and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO ₂ and pH given
	flag = 22 pCO ₂ and HCO ₃ given
	flag = 23 pCO ₂ and CO ₃ given
	flag = 24 pCO ₂ and ALK given
	flag = 25 pCO ₂ and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO ₂ in μatm
var2	Value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)

b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

This subroutine has same input parameters as subroutine carb(). For details on these parameters, refer to documentation of 'carb'.

This subroutine computes partial derivatives of each output variable with respect to each of the input variable (including each of the two chosen carbonate system variables, each of the nutrients (total silicon and total phosphorus), temperature, and salinity).

It computes these derivatives (dy/dx) using the method of central differences, i.e.,

- for dx, it adds a positive and negative perturbation, same and equal in magnitude, to each input variable, one at a time, and
- for dy, it then computes the corresponding induced change in output variables

All arguments but the first (varid), can be given as scalars or vectors. If the lengths of the vectors differs, only the longest vector is retained and the other arguments are set equal to the first value of the other vectors are used. Hence users should use either vectors with the same dimension or one vector for one argument and scalars for others; otherwise, results may not be as intended.

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

H	derivative of [H+] concentration (mol/kg/...)
pH	derivative of pH

CO2	derivative of CO2 concentration (mol/kg/...)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure ($\mu\text{atm}/...$)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure ($\mu\text{atm}/...$)
HCO3	derivative of HCO3 concentration (mol/kg/...)
CO3	derivative of CO3 concentration (mol/kg/...)
DIC	derivative of DIC concentration (mol/kg/...)
ALK	derivative of ALK, total alkalinity (mol/kg/...)
OmegaAragonite	derivative of Omega aragonite, aragonite saturation state
OmegaCalcite	derivative of Omega calcite, calcite saturation state

If all input data have the same 'flag' value, returned data frame does not show derivatives of input pair of carbonate system variables. For example, if all input flags are 15, the input pair is DIC and ALK; hence, derivatives of DIC and ALK are not returned.

Units of derivative dy/dx is unit(y)/unit(x) where unit(x) are as follows:

degree C	when x is Temperature
psu	when x is Salinity
μatm	when x is pCO2
mol/kg	for all other cases

Author(s)

Jean-Marie Epitalon, James Orr, and Jean-Pierre Gattuso<gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **22**, vol49 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO2 calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHE: 149 (2013) 8-22], *Marine Chemistry*, **165**, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```

## 1) For the input pair ALK and DIC (var1 and var2 when flag=15)
##   compute derivatives of all output varialbes
##   with respect to DIC (i.e., var2)
derivnum(varid='var2', flag=15, var1=2300e-6, var2=2000e-6,
          S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
          pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## 2) For the input pair pH and ALK (var1 and var2 when flag=8)
##   compute derivatives of all output variables
##   with respect to [H+] concentration
derivnum(varid='var1', flag=8, var1=8.2, var2=0.00234,
          S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
          pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## 3) Using vectors as arguments and compute derivatives of all output
##   variables with respect to temperature
flag <- c(8, 2, 8)

```

```

var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("1", "1", "1")
pHscale <- c("T", "T", "T")
b <- c("u74", "u74", "u74")
derivnum(varid='T', flag=flag, var1=var1, var2=var2, S=S, T=T, P=P,
         Pt=Pt, Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

# For more examples of use of derivnum.R,
# consult the code of seacarb's errors routine.

```

eos2teos_chem

Convert temperature and salinity from EOS-80 to TEOS-10

Description

Converts in situ temperature to conservative temperature and practical to absolute salinity (SA). Salinity conversion depends on total alkalinity as well as the concentrations of dissolved inorganic carbon, nitrate and silicate.

Usage

```
 eos2teos_chem(SP, T, P=0, TA=2300e-6, DIC=2000e-6, NO3=0, SIOH4=0)
```

Arguments

SP	Practical salinity on the practical salinity scale
T	In situ temperature in deg. C
P	Sea water pressure in dbar
TA	Total alkalinity, in mol/kg, default is 2300 $\mu\text{mol/kg}$
DIC	Dissolved inorganic carbon concentration in mol/kg, default is 2000 $\mu\text{mol/kg}$
NO3	Total nitrate concentration in mol/kg, default is 0
SIOH4	Total silicate concentration in mol/kg, default is 0

Details

Conversion from practical to absolute salinity depends on carbonate system parameters and ion concentration which mostly affect water density anomalies.

Value

The function returns a data frame containing the following columns:

CT	Conservative temperature (deg C)
SA	Absolute salinity (g/kg)

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., Wright D. G. and Millero F. J., 2011. The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* **7**, 363-387.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

teos2eos_chem does the reverse, eos2teos_geo, sp2sa_chem
package gsw

Examples

```
# Calculate Conservative Temperature and Absolute Salinity of a sample with
# Practical salinity of 35 psu, in-situ temperature of 18 deg C,
# at 0 dbar and total alkalinity of 0.00234 mol/kg and DIC of 0.00202 mol/kg
f <- eos2teos_chem(SP=35, T=18, P=0, TA=0.00234, DIC=0.00202)
CT <- f$CT      # Conservative Temperature
SA <- f$SA      # Absolute Salinity
```

eos2teos_geo

Convert temperature and salinity from EOS-80 to TEOS-10

Description

Converts in situ to conservative temperature and practical to absolute salinity (SA). Salinity conversion depends on depth and geographic location.

Usage

```
 eos2teos_geo(SP, T, P=0, long=1.e20, lat=1.e20)
```

Arguments

SP	Practical salinity on the practical salinity scale
T	In situ temperature in deg. C
P	Sea water pressure in dbar
long	Longitude in decimal degrees [0 ... +360] or [-180 ... +180]
lat	Latitude in decimal degrees [-90 ... 90]

Details

Conversion from practical to absolute salinity depends on water density anomaly which is correlated with silicate concentration. This function relies on silicate concentration taken from WOA (World Ocean Atlas) to evaluate density anomaly.

Value

The function returns a data frame containing the following columns:

CT	Conservative temperature (deg C)
SA	Absolute salinity (g/kg)

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

teos2eos_geo does the reverse, eos2eos_chem, sp2sa_geo, package gsw

Examples

```
# Calculate conservative temperature and absolute salinity of a sample with
# Practical salinity of 35 psu, in situ temperature of 18 deg C,
# depth is 10 dbar and location is 188 degrees East and 4 degrees North.
f <- eos2eos_geo(SP=35, T=18, P=10, long=188, lat=4)
CT <- f$CT      # Conservative temperature
SA <- f$SA      # Absolute salinity
```

errors*Uncertainty propagation for computed marine carbonate system variables*

Description

Estimates combined standard uncertainties in computed carbonate system variables by propagating input uncertainties (standard uncertainties) in six input variables, including (Orr et al., Mar. Chem., in press):

- the input pair of carbonate system variables,
- the 2 input nutrients (silicate and phosphate concentrations),
- temperature and salinity. It also accounts for
- the errors in the key dissociation constants pK_0 , pK_1 , pK_2 , pK_b , pK_w , pK_{spa} and pK_{spc}
- the error in total boron

Usage

```
errors(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
       evar1=0, evar2=0, eS=0.01, eT=0.01, ePt=0, eSit=0,
       epK=c(0.002, 0.0075, 0.015, 0.01, 0.01, 0.02, 0.02),
       eBt=0.02, method = "ga", r=0.0, runs=10000,
       k1k2='x', kf='x', ks="d", pHscale="T", b="u74", gas="potential",
       warn="y", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the pair of carbonate system input variables. The flags to be used are as follows: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given flag = 6 pH and HCO ₃ given flag = 7 pH and CO ₃ given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO ₃ and CO ₃ given flag = 11 HCO ₃ and ALK given flag = 12 HCO ₃ and DIC given flag = 13 CO ₃ and ALK given flag = 14 CO ₃ and DIC given flag = 15 ALK and DIC given
------	---

	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	Value of the first variable (in mol/kg, except for pH and for pCO2 in μ atm)
var2	Value of the second variable (in mol/kg, except for pH)
S	Salinity (practical salinity scale)
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total dissolved inorganic phosphorus (mol/kg); set to 0 if NA
Sit	Concentration of total dissolved inorganic silicon (mol/kg); set to 0 if NA
evar1	Standard uncertainty in var1 of input pair of carbonate system variables
evar2	Standard uncertainty in var2 of input pair of carbonate system variables
eS	Standard uncertainty in salinity; default is 0.01
eT	Standard uncertainty in temperature (degree C); default is 0.01
ePt	Standard uncertainty in total dissolved inorganic phosphorus concentration (mol/kg)
eSit	Standard uncertainty in total dissolved inorganic silicon concentration (mol/kg)
epK	Standard uncertainty) in 7 key dissociation constants: pK0, pK1, pK2, pKb, pKw, pKspa and pKspc. This is a vector. The default is c(0.002, 0.0075, 0.015, 0.01, 0.01, 0.02, 0.02).
eBt	Standard uncertainty in total boron, given as a relative fractional error. The default is 0.02, which equates to a 2% error
method	Case insensitive character string to choose the error-propagation method: 1) Gaussian, 2) Method of Moments, or 3) Monte Carlo). These methods are specified using the 2-letter codes "ga", "mo", or "mc", respectively. The default is "ga" (Gaussian).
r	Correlation coefficient between standard uncertainties of var1 and var2 (only useful with method="mo", i.e., ignored for the 2 other methods, the default is r=0.0)
runs	Number of random samples (ignored unless method="mc"; the default is runs=10000
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".

ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

Complete information on routine uncertainty propagation for the marine carbon dioxide system can be found in Orr et al. (in press). This function requires users to specify each input standard uncertainty as either the standard deviation or the standard error of the mean. The latter implies much smaller propagated uncertainties, but is appropriate only when interested in the error in the mean, not the error of a given measurement. Beware that it is easy to fool oneself when using the standard error of the mean rather than the standard deviation.

This function requires different types of standard uncertainties:

- Standard uncertainties for evar1, evar2, eS, eT, ePt, eSit (same units as the input data, e.g., mol/kg);
- Standard uncertainties in pK units for epK; and
- Standard uncertainties in relative fractional units (between 0.0 and 1.0) for eBt.

This function propagates standard uncertainty from input to output variables using one of three methods:

- Gaussian: The Gaussian method is the standard technique for estimating a computed variable's (z) second moment (its variance or standard deviation) based on a first-order approximation to z. More precisely, we use here the basic 1st order, 2nd moment uncertainty analysis (a type of Taylor expansion), assuming no covariance between input variables. This is the approach used by Dickson and Riley (1978). It is the default method.
- Method of moments: The method of moments is a more general form of the Gaussian method. But in addition, it also accounts for covariance between input variables. In this case, the

'errors' routine allows the user to specify a value of the correlation coefficient 'r', having a value between -1.0 and 1.0, to indicate the correlation between standard uncertainties of the input pair of carbonate system variables. That correlation is used to compute the covariance. But by default, it is assumed that there is no covariance ($r=0.0$).

- Monte Carlo: The Monte Carlo method is a brute-force approach relying on repeated random sampling of input errors, adding those to each input variables, calculating the corresponding output variables for each sample, and finally assessing the standard deviation in each output variables.

This function has many input parameters that are identical to those in the carb function. For their details, refer to the 'carb' documentation.

All parameters may be scalars or vectors except epK, eBt, method, runs, and gas.

- runs and eBt must be scalars
- method and gas must each consist of a character string
- epK may be a vector of 7 values. In that case, it must list errors for pK0, pK1, pK2, pKb, pKw, pKspa and pKspc, respectively. That set of errors is identical for all input data. Alternatively, users may specify 'epK=NULL' or 'epK=0' to set all 7 values to zero and thus neglect errors in the equilibrium constants.

In contrast, for evar1, evar2, r, eS, eT, ePt and eSit:

- if they are vectors, they represent standard uncertainties associated with each data point
- if they are scalars (single real numbers), they represent one standard uncertainty value each associated to all data points

The same remark applies to parameter r (correlation coefficient).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a 2-dimensional dataframe, with the following columns:

- H combined standard uncertainty in $[H^+]$ concentration (mol/kg)
- pH combined standard uncertainty in pH
- CO2 combined standard uncertainty in CO2 concentration (mol/kg)
- pCO2combined standard uncertainty in "standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μatm)
- fCO2combined standard uncertainty in "standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μatm)
- HCO3combined standard uncertainty in HCO3 concentration (mol/kg)

- CO3combined standard uncertainty in CO3 concentration (mol/kg)
- DICcombined standard uncertainty in DIC concentration (mol/kg)
- ALKcombined standard uncertainty in ALK, total alkalinity (mol/kg)
- OmegaAragonitecombined standard uncertainty in Omega aragonite (aragonite saturation state)
- OmegaCalcitecombined standard uncertainty in Omega calcite (calcite saturation state)

If all input data have the same 'flag' value, the returned data frame does not show combined standard uncertainties on input pair of carbonate system variables. For example, if all input flags are 15, the input pair is DIC and ALK; hence, errors on DIC and ALK are not returned.

Correlation coefficient

By default, 'r' is zero. However, for some pairs the user may want to specify a different value. For example, measurements of pCO2 and pH are often anti-correlated. The same goes for two other pairs: 'CO2 and CO3' and 'pCO2 and CO3'. But even for these cases, care is needed before using non-zero values of 'r'.

When the user wishes to propagate standard uncertainties for an individual measurement, 'r' should ALWAYS be zero if each member of the input pair is measured independently. In this case, we are interested in the correlation between the uncertainties in those measurements, not in the correlation between the measurements themselves. Uncertainties from those measurements are probably not correlated if they come from different instruments. Conversely, if users are interested in the error in the mean of a distribution of measurements (i.e., if they are propagating standard errors instead of standard deviations), one should then also account for the correlation between the measurements of the two variables of the input pair.

For input pairs where one member is pH (flags 1, 6, 7, 8, 9, and 21), this 'errors' function automatically inveres the sign of 'r'. The reason for that is that the associated derivatives are computed in terms of the hydrogen ion concentration (H+), not pH. Therefore for each of these 6 flags, if the user wants to compute their own 'r' that should be done by (1) using the H+ concentration instead of pH, and (2) invering the sign of that computed 'r' before passing it as an argument to this routine. Usually though (when not calculating r for pH), the user may just use the 'r' in the expected way. For example, to include the covariance term when there is a perfect anticorrelation of pH with pCO2, one would use 'r=-1.0'.

Computation time

Computation time depends on the method chosen; the Monte Carlo method takes much longer to execute. The computational time required for the Monte Carlo method is proportional to the number of runs. More runs, implies improved accuracy: runs = 10000 appears a minimum to obtain an accuracy of less than 1%. Accuracy is inversely proportional to the number of runs.

Computation time also depends on the chosen pair of input variables. For example, with the input pair DIC and Total alkalinity (flag=15), it is much longer than for input pair pH and Total alkalinity (flag=8)

Author(s)

Jean-Marie Epitalon, James Orr, and Jean-Pierre Gattuso<gattuso@obs-vlfr.fr>

References

Dickson, A. G. and Riley, J. P., 1978 The effect of analytical error on the evaluation of the components of the aquatic carbon-dioxide system, *Marine Chemistry*, **6**, 77-85.

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **22**, vol49 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M., Dickson A. and Gattuso J.-P., in press. Routine uncertainty propagation for the marine carbon dioxide system. *Marine Chemistry*.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry*, **165**, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```

## 1) For the input pair ALK and DIC (var1 and var2 when flag=15),
## compute resulting uncertainty from given uncertainty on ALK and DIC (5 umol/kg)
## and default uncertainties in dissociation constants and total boron
## using the default method (Gaussian)
errors(flag=15, var1=2300e-6, var2=2000e-6, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
      evar1=5e-6, evar2=5e-6, eS=0, eT=0, ePt=0, eSit=0,
      pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
## Typical output:
## H          pH          CO2          fCO2          pCO2          HCO3      ...
## 3.721614e-10  0.01796767  5.441869e-07  19.25338  19.31504  9.170116e-06  ...
## 3.721614e-10  0.01796767  5.441869e-07  19.25338  19.31504  9.170116e-06  ...

## 2) Do the same as in one, but assign a 4% uncertainty to total boron
## This uncertainty is the amount by which estimates from Lee et al (2010) and
## Uppstrom (1974) differ. The default for the latter is eBt=0.02.
## errors(flag=15, var1=2300e-6, var2=2000e-6, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
##        evar1=5e-6, evar2=5e-6, eS=0, eT=0, ePt=0, eSit=0, eBt=0.04,
##        pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## 3) For the input pair pH and ALK (var1 and var2 when flag=8)
## compute standard errors in output variables from errors in input variables, i.e.,
## for pH (0.005 pH units) and in ALK (5 umol/kg), along with
## errors in total dissolved inorganic phosphorus (0.1 umol/kg) and
## total dissolved inorganic silicon (2 umol/kg) concentrations, while
## assuming no uncertainty in dissociation constants & boron, using the Gaussian method:
## errors(flag=8, var1=8.25, var2=2300e-6, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
##        evar1=0.005, evar2=5e-6, eS=0, eT=0, ePt=0.1, eSit=2, epK=0, eBt=0,
##        method="ga", pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## 4) For the input pair pCO2 and pH (var1 and var2 when flag=21)
## compute standard errors in output variables from errors in input variables, i.e.,
## for pCO2 (2 uatm) and pH (0.005 pH units), with no uncertainties in Pt and Sit
## nor in the dissociation constants BUT a perfect anticorrelation between pCO2 and pH,
## (the input pair) using the Method of moments:
## errors(flag=21, var1=400, var2=8.1, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
##        evar1=2, evar2=0.005, eS=0, eT=0, ePt=0.0, eSit=0, epK=0, eBt=0,
##        method="mo", r=-1.0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## 5) Use vectors as arguments and compute errors on all output variables
## using Monte Carlo method taking into account input errors on pH, ALK, DIC
## and dissociation constants (pKx)
flag <- c(8, 15)
var1 <- c(8.2, 0.002394, 8.25)

```

```

var2 <- c(0.002343955, 0.002017)
S <- c(35, 35)
T <- c(25, 25)
P <- 0
Pt <- 0
Sit <- 0
evar1 <- c(0.005, 2e-6)
evar2 <- c(2e-6, 2e-6)
epKx <- c(0.002, 0.01, 0.02, 0.01, 0.01, 0.01, 0.01)
eBtx = 0.01
method <- "mc"
kf <- "pf"
k1k2 <- "1"
pHscale <- "T"
b <- "u74"
## NOTE that the following is executable but enclosed in "donttest"
## because it takes too long to run when submiting to CRAN
## and is therefore rejected
errors(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt, Sit=Sit,
       evar1=evar1, evar2=evar2, eS=0, eT=0, ePt=0, eSit=0, epK=epKx, eBt=eBtx,
       method=method, runs=10000, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

```

f2pCO2*Converts the CO2 fugacity to CO2 partial pressure*

Description

Converts fCO2 (fugacity of CO2) into pCO2 (partial pressure in CO2)

Usage

```
f2pCO2(T = 25, Patm=1, P=0, fCO2)
```

Arguments

T	Temperature in degrees Celsius, default is 25oC
Patm	Surface atmospheric pressure in atm, default 1 atm
P	Hydrostatic pressure in bar, default is 0 bar (surface)
fCO2	Fugacity of CO2 in μ atm, the same units as that for the pCO2 output

Value

pCO2	Partial pressure of CO2 in μ atm, the same units as that for the fCO2 input
------	---

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Heloise Lavigne, Jean-Pierre Gattuso, and James Orr <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007. Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359, 1980.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

See Also

[p2fCO2](#).

Examples

```
f2pCO2(T=25, Patm=1.0, P=0, fCO2=380)
```

fCO2insi

fCO2 at in situ temperature

Description

Correction to compensate for the difference in temperature between the temperature of measurement and in situ temperature.

Usage

```
fCO2insi(fCO2lab = 400, Tlab = 20, SST = 19)
```

Arguments

fCO2lab	Fugacity of CO ₂ measured in the lab in μatm
Tlab	Temperaure of measurment in the lab in in degrees Celsius, default is 20oC
SST	Temperature in degrees Celsius, default is 19oC

Value

fCO2insi	Fugacity of CO ₂ at in situ temperature in μatm , the same units as that for the fCO ₂ input.
----------	--

Note

The empirical correction applied is from Takahashi (1993) as recommended by Pierrot et al. (2009)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Pierrot D., Neill C., Sullivan K., Castle R., Wanninkhof R., LÃ¼ger H., Johannessen T., Olsen A., Feely R. A. and Cosca C. E., 2009. Recommendations for autonomous underway pCO₂ measuring systems and data-reduction routines. *Deep-Sea Res. II* **56**, 512-522.

Takahashi T., Olafsson J., Goddard J. G., Chipman D. W. and Sutherland S. C., 1993. Seasonal variation of CO₂ and nutrients in the high-latitude surface oceans: a comparative study. *Glob. Biogeochem. Cycles* **7**, 843-878.

See Also

[f2pCO2](#), [pCO2insi](#).

Examples

```
fCO2insi(fCO2lab = 400, SST = 15, Tlab = 16)
```

K0

Henry's constant mol/(kg/atm)

Description

Henry's constant mol/(kg/atm)

Usage

```
K0(S=35, T=25, P=0, Patm=1, warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
Patm	Surface atmospheric pressure in atm, default is 1 atm
warn	"y" to show warnings when T or S go beyond the valid range for K0; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between -1 and 45oC.

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

For pressure corrections: the pressure correction term of Weiss (1974) is used.

Value

K0 Henry's constant mol/(kg/atm)

Author(s)

Jean-Marie Epitalon, Aurelien Proye, and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Weiss R. F., 1974 Carbon dioxide in water and seawater: the solubility of a non-ideal gas. *Marine Chemistry* **2**, 203-215.

Examples

K0(S=35, T=25, P=0)

K1

First dissociation constant of carbonic acid (mol/kg)

Description

First dissociation constant of carbonic acid (mol/kg)

Usage

K1(S=35, T=25, P=0, k1k2="x", pHscale="T", kSWS2scale="x", ktotal2SWS_P0="x", warn="y")

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0

k1k2	"I" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "I", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is not required for all formulations of K1 and K2, nor when the hydrostatic pressure is 0. It is advised to use default value "x", in which case it is computed when required.
ktotal2SWS_P0	Conversion factor from the total scale to the SWS at an hydrostatic pressure of 0. It is not required for all formulations of K1 and K2. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for K1; "n" to suppress warnings. The default is "y".

Details

The Lueker et al. (2000) constant is recommended by Guide to Best Practices for Ocean CO₂ Measurements (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): formulation is that of Waters et al (2014). See below.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

The pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

K1	First dissociation constant of carbonic acid (mol/kg)
----	---

Author(s)

Jean-Marie Epitalon and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Lueker T. J., Dickson A. G., and Keeling C. D., 2000 Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., Pierrot D., Lee K., Wanninkhof R., Feely R., Sabine C. L., Key R. M. and Takahashi T., 2002. Dissociation constants for carbonic acid determined from field measurements. Deep Sea Research Part I: Oceanographic Research Papers **49**:1705-1723.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H., and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Mojica Prieto F. J. and Millero F. J., 2002. The values of pK1 + pK2 for the dissociation of carbonic acid in seawater. *Geochimica et Cosmochimica Acta* **66**, 2529-2540.

Papadimitriou S., Loucaides S., RÃ©rolle V. M. C., Kennedy P., Achterberg E. P., Dickson A. G., Mowlem M. and Kennedy H., 2018. The stoichiometric dissociation constants of carbonic acid in seawater brines from 298 to 267 K. *Geochimica et Cosmochimica Acta* **220**, 55-70.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*..

Sulpis O., Lauvset S. K. and Hagens M., 2020. Current estimates of K1* and K2* appear inconsistent with measured CO₂ system parameters in cold oceanic regions. *Ocean Science* **16**, 847-862.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHE: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

See Also[K2.](#)**Examples**

```
K1(S=35, T=25, P=0, k1k2="1", pHscale="T")
```

K1p*First dissociation constant of phosphoric acid (mol/kg)*

Description

First dissociation constant of phosphoric acid (mol/kg)

Usage

```
K1p(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	Choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for K1p; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45oC.

The pressure correction was applied on the seawater scale. Hence, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

```
K1p
```

First dissociation constant of phosphoric acid (mol/kg)

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

See Also

[K2p](#), [K3p](#).

Examples

`K1p(35, 25, 0)`

K2	<i>Second dissociation constant of carbonic acid (mol/kg)</i>
----	---

Description

Second dissociation constant of carbonic acid (mol/kg)

Usage

`K2(S=35, T=25, P=0, k1k2="x", pHscale="T", kSWS2scale="x", ktotal2SWS_P0="x", warn="y")`

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
k1k2	"I" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "I", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale

kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is not required for all formulations of K1 and K2, nor when the hydrostatic pressure is 0. It is advised to use default value "x", in which case it is computed when required.
ktotal2SWS_P0	Conversion factor from the total scale to the SWS at an hydrostatic pressure of 0. It is not required for all formulations of K1 and K2. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for K2; "n" to suppress warnings. The default is "y".

Details

The Lueker et al. (2000) constant is recommended by Guide to Best Practices for Ocean CO₂ Measurements (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): S ranging from 19.6 to 41 and T ranging between 15 to 35oC.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

The pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

K2	Second dissociation constant of carbonic acid (mol/kg)
----	--

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

The Lueker et al. (2000) constant is recommended by Guide to Best Practices for Ocean CO₂ Measurements (2007). The Roy et al. (1993) constants is recommended by DOE (1994).

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Mojica Prieto F. J. and Millero F. J., 2002. The values of pK1 + pK2 for the dissociation of carbonic acid in seawater. *Geochimica et Cosmochimica Acta* **66**, 2529-2540.

Papadimitriou S., Loucaides S., RÃ©rolle V. M. C., Kennedy P., Achterberg E. P., Dickson A. G., Mowlem M. and Kennedy H., 2018. The stoichiometric dissociation constants of carbonic acid in seawater brines from 298 to 267 K. *Geochimica et Cosmochimica Acta* **220**, 55-70.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Sulpis O., Lauvset S. K. and Hagens M., 2020. Current estimates of K1* and K2* appear inconsistent with measured CO₂ system parameters in cold oceanic regions. *Ocean Science* **16**, 847-862.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHE: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

See Also

[K1](#).

Examples

$\text{K2}(35, 25, 0)$

K2p	<i>Second dissociation constant of phosphoric acid (mol/kg)</i>
-----	---

Description

Second dissociation constant of phosphoric acid (mol/kg)

Usage

```
K2p(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for K2p; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45oC.

The pressure correction was applied on the seawater scale. Hence, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

K2p Second dissociation constant of phosphoric acid (mol/kg)

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

See Also

[K1p](#), [K3p](#).

Examples

`K2p(35,25,0)`

K2si

Second dissociation constant of Si(OH)4

Description

Second dissociation constant of Si(OH)4 (mol/kg)

Usage

`K2si(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", ktotal2SWS_P0="x")`

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
ktotal2SWS_P0	Conversion factor from the total scale to the SWS at an hydrostatic pressure of 0. It is advised to use default value "x", in which case it is computed when required.

Details

This equation is modified from Wischmeyer et al. (2003), who fitted the temperature-dependent K2si from Nordstrom et al. (1990) for freshwater to a value of 12.56 for T=25 and an ionic strength of 0.5 mol/kg. The temperature and salinity ranges in which it is valid are not well constrained.

The pressure correction is applied on the seawater scale. Hence, values are first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value is transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

K2si Second dissociation constant of Si(OH)4 (mol/kg)

Author(s)

Mathilde Hagens (<M.Hagens@uu.nl>)

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Nordstrom D. K., L. N. Plummer, D. Langmuir, E. Busenberg, H. M. May, B. F. Jones, D. L. Parkhurst, 1990 Revised chemical equilibrium data from major mineral reactions and their limitations. In: Melchior, D.C., R. L. Bassett (Eds.) Chemical Modeling of Aqueous Systems. *IIACS Series* **416**. American Chemical Society, Washington, DC.

Wischmeyer A. G., Y. Del Amo, M. Brzezinski, D. A. Wolf-Gladrow, 1995 Theoretical constraints on the uptake of silicic acid species by marine diatoms. *Marine Chemistry* **82**: 13-29.

Examples

```
K2si(S=35, T=25, P=0, pHscale="T")
```

K3p

Third dissociation constant of phosphoric acid (mol/kg)

Description

Third dissociation constant of phosphoric acid (mol/kg)

Usage

```
K3p(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for K3p; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45oC.

The pressure correction was applied on the seawater scale. Hence, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

K3p	Third dissociation constant of phosphoric acid (mol/kg)
-----	---

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication 3*, 1-191.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

See Also

[K1p](#), [K2p](#).

Examples

`K3p(35, 25, 0)`

K_b *Dissociation constant of boric acid (mol/kg)*

Description

Dissociation constant of boric acid (mol/kg)

Usage

```
Kb(S=35, T=25, P=0, pHscale="T", ksWS2scale="x", ktotal2SWS_P0="x", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
ktotal2SWS_P0	Conversion factor from the total scale to the SWS at an hydrostatic pressure of 0. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for Kb; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 5 and 45 and T ranging between 0 and 45°C.

The pressure correction is applied on the seawater scale. Hence, values are first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

K_b Dissociation constant of boric acid (mol/kg)

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Prove and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Thermodynamics of the dissociation of boric acid in synthetic seawater from 273.15 to 318.15 K. *Deep-Sea Research* **375**, 755-766.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Examples

```
Kb(S=35, T=25, P=0, pHscale="T")
```

kconv

Conversion factors to change the pH scale of dissociation constants

Description

Conversion factors from the total scale to the free and seawater scales

Usage

```
kconv(S=35, T=25, P=0, kf, Ks, Kff, warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994); if the fonction Kf was used previously, the default value is the value given for the argument kf in the fonction Kf. If the Kf function was not used previously, the default value is "pf", except if T is outside the range 9 to 33oC or of S is outside the range 10 to 40. In these cases, the default is "dg".
Ks	Stability constant of hydrogen sulfate (mol/kg) at given S, T and P, optional; if not given, it will be computed, if given, it allows for speed optimisation
Kff	Stability constant of hydrogen fluoride (mol/kg) on free pH scale at given S, T and P, optional; if not given, it will be computed, if given, it allows for speed optimisation and kf parameter is then ignored
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to supress warnings. The default is "y".

Details

It is critical to consider that each formulation is valid in specific ranges of temperature and salinity:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

Note that kconv may be called in many functions (i.e. K1, K2, K1p, K2p, K3p, Kw, Ks_i, K2s_i, etc...) without user controls it. To force a particular formulation for K_f, it is recommended to call kconv() first then pass the resulting conversion factors to these functions.

Value

The function returns a list with 6 conversion factors :

ktotal2SWS	to convert from the total scale to seawater scale
ktotal2free	to convert from the total scale to the free scale
kfree2SWS	to convert from the free scale to the seawater scale
kfree2total	to convert from the free scale to total scale
kSWS2total	to convert from the seawater scale to the total scale
kSWS2free	to convert from the seawater scale to the free scale

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl>

References

Dickson A.G. and F.J. Millero, 1987 A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Research* **34**:1733-1743.

See Also

[pHconv.](#)

Examples

```
##To convert dissociation constants from the total scale to the free scale
## (at salinity=35, temperature=25oC and atmospheric pressure):
kconv(35,25,0)
conv <- kconv()
c(K1_total=K1(),K1_SWS=K1()*conv$ktotal2SWS,K1_free=K1()*conv$ktotal2free)
```

Kf*Equilibrium constant of hydrogen fluoride (mol/kg)*

Description

Stability constant of hydrogen fluoride (mol/kg)

Usage`Kf(S=35, T=25, P=0, kf="x", pHscale="T", Ks_p0="x", Ks_p="x", warn="y")`**Arguments**

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
kf	"pf" for using Kf from Perez and Fraga (1987) "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994), default is "pf". Attention do not use a vector for this argument.
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
Ks_p0	Stability constant of hydrogen sulfate (mol/kg) at pressure zero; needed if kf = "pf"; if needed and not given or set to "x", it is computed; if given, computation speed is increased
Ks_p	Stability constant of hydrogen sulfate (mol/kg) at chosen pressure; if not given or set to "x", it is computed; if given, computation speed is increased
warn	"y" to show warnings when T or S go beyond the valid range for Kf; "n" to suppress warnings. The default is "y".

Details

The Perez and Fraga (1987) constant is recommended by Guide to Best Practices for Ocean CO₂ Measurements (2007). The Dickson and Riley (1979 in Dickson and Goyet, 1994) constant is recommended by DOE (1994).

It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

The pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

Kf Stability constant of hydrogen fluoride (mol/kg)

Author(s)

Jean-Marie Epitalon, Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Examples

```
Kf(S=35, T=25, P=0, kf="pf", pHscale="T")
```

kfg *variable for internal use*

Description

nothing

Khs	<i>Dissociation constant of hydrogen sulfide (mol/kg)</i>
-----	---

Description

Dissociation constant of hydrogen sulfide (mol/kg)

Usage

```
Khs(S=35, T=25, P=0, pHscale="T", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
warn	"y" to show warnings when T or S go beyond the valid range for Khs; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45oC.

The pressure correction is applied on the seawater scale. Hence, the values are first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value is transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Khs Dissociation constant of hydrogen sulfide

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl> and Heloise Lavigne

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* 59: 661-677.

Examples

```
Khs(S=35,T=25,P=0, pHscale="T")
```

Kn *Dissociation constant of ammonium (mol/kg)*

Description

Dissociation constant of ammonium on the total scale (mol/kg)

Usage

```
Kn(S=35, T=25, P=0, pHscale="T", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
warn	"y" to show warnings when T or S go beyond the valid range for Kn; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45°C.

The pressure correction is applied on the seawater scale. Hence, values are first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value is transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Kn Dissociation constant of ammonium (mol/kg)

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl> and Heloise Lavigne

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Examples

```
Kn(S=35, T=25, P=0, pHscale="T")
```

Ks	Stability constant of hydrogen sulfate (mol/kg)
----	---

Description

Stability constant of hydrogen sulfate (mol/kg)

Usage

```
Ks(S=35, T=25, P=0, ks="d", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
warn	"y" to show warnings when T or S go beyond the valid range for Ks; "n" to suppress warnings. The default is "y".

Details

The Dickson (1990) constant is recommended by Guide to Best Practices for Ocean CO₂ Measurements (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

The pressure correction is applied on the free scale as described by Millero (1995), and the value transformed back to the required scale (T, F or SWS).

Value

Ks	Stability constant of hydrogen sulfate (mol/kg), pHscale = free scale
----	---

Author(s)

Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Examples

```
Ks(S=35, T=25, P=0, ks="d")
```

Ksi*Dissociation constant of Si(OH)_4*

Description

Dissociation constant of Si(OH)_4 on total scale (mol/kg)

Usage

```
Ksi(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25°C
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for Ks; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45°C.

The pressure correction is applied on the seawater scale. Hence, values are first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value is transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl> and Heloise Lavigne

References

DOE 1994 *Handbook of methods for the analysis of the various parameters of the carbon dioxide system in sea water*. ORNL/CDIAC-74. Oak Ridge, Tenn.: Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Examples

```
Ksi(S=35, T=25, P=0, pHscale="T")
```

Kspa

Usage

KoneCS-2E-T-2E-B-2 (wapp="1111")

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
warn	"y" to show warnings when T or S go beyond the valid range for Kspa; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 5 and 44 and T ranging between 5 and 40oC.

Pressure correction was performed as described by Millero (1995).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Kspa Solubility product of aragonite (mol²/kg)

Author(s)

Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59** 661-677.

Mucci A., 1983 The solubility of calcite and aragonite in seawater at various salinities, temperature, and one atmosphere total pressure. *American Journal of Science* **283**: 780-799.

See Also

[Kspc](#).

Examples

Kspa(S=35, T=25, P=0)

Kspc	<i>Solubility product of calcite (mol/kg)</i>
------	---

Description

Solubility product of calcite (mol/kg)

Usage

```
Kspc(S=35, T=25, P=0, warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
warn	"y" to show warnings when T or S go beyond the valid range for Kspc; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 5 and 44 and T ranging between 5 and 40oC.

The pressure correction was performed as described by Millero (1995).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Kspc Solubility product of calcite (mol2/kg)

Author(s)

Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59** 661-677.

Mucci A., 1983 The solubility of calcite and aragonite in seawater at various salinities, temperature, and one atmosphere total pressure. *American Journal of Science* **283**: 780-799.

See Also[Kspa](#).**Examples**`Kspc(S=35, T=25, P=0)`

Kw	<i>Ion product of water (mol2/kg2)</i>
----	--

Description

Ion product of water (mol2/kg2)

Usage
`Kw(S=35, T=25, P=0, pHscale="T", kSWS2scale="x", warn="y")`
Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
kSWS2scale	Conversion factor from the seawater scale (SWS) to the pH scale selected at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required.
warn	"y" to show warnings when T or S go beyond the valid range for Kw; "n" to suppress warnings. The default is "y".

Details

This formulation is only valid for specific ranges of temperature and salinity:

- S ranging between 0 and 45 and T ranging between 0 and 45oC.

The pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

Kw	Ion product of water (mol2/kg2)
----	---------------------------------

Author(s)

Heloise Lavigne, Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59** 661-677.

Examples

```
Kw(S=35, T=25, P=0, pHscale="T")
```

oa

Perturbation of the seawater carbonate system

Description

Describes the various approaches that can be used to alter the seawater carbonate system. Its main purpose is to assist the design of ocean acidification perturbation experiments.

Usage

```
oa(flag, var1, var2, pCO2f, pCO2s=1e6, S=35, T=25, P=0,
Pt=0, Sit=0, k1k2='x', kf='x', ks="d", pHscale="T", plot=FALSE,
b="u74", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available to describe the initial seawater. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given flag = 6 pH and HCO ₃ given flag = 7 pH and CO ₃ given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO ₃ and CO ₃ given flag = 11 HCO ₃ and ALK given flag = 12 HCO ₃ and DIC given flag = 13 CO ₃ and ALK given
------	---

	flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
var1	Value of the first variable available to describe the initial seawater, in mol/kg except for pH and for pCO2 in uatm
var2	Value of the second variable available to describe the initial seawater, in mol/kg except for pH
pCO2f	pCO2 target value, in uatm
pCO2s	pCO2s is the pCO2, in uatm, of the "high-CO2" seawater that will be mixed with "normal seawater". The default value is 10^6 uatm, that is seawater bubbled with pure CO2 gas and saturated with CO2.
S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25
P	Hydrostatic pressure in bar (surface = 0), default is 0
Pt	Concentration of total phosphate in mol/kg, default is 0
Sit	Concentration of total silicate in mol/kg, default is 0
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
plot	A plot of the different perturbation methods can be plotted in a DIC vs ALK field with pCO2 isolines are drawn in the back. Default is false.
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_f , the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

The function returns a list built as follows:

description	A table describing in plain English the various ways to reach the target pCO ₂ . Note that if a vector is given in argument only the first value is used.
perturbation	Table providing key parameters for the following methods: <i>CO₂ bubbling</i> : high-CO ₂ air is bubbled in seawater. The first parameter is the value of the pCO ₂ in the air required to bubble the seawater (in uatm). <i>SW mixing</i> : mixing of “normal” and “high-CO ₂ ” seawater. The first parameter, “Weight fraction high-CO ₂ SW” or w_f , is the weight fraction of the high-CO ₂ seawater per kg seawater. <i>Addition of acid</i> : strong acid is added to seawater. Note that this method is not recommended because it does not closely mimic natural ocean acidification (Gattuso and Lavigne, 2009). The first parameter, H ⁺ (mol/kg), is the amount of H ⁺ that must be added (mol/kg). The acid must be fortified with NaCl in order to have the same salinity than seawater. <i>Addition of HCO₃ and acid</i> : bicarbonate (HCO ₃) and a strong acid are added. The first parameter, HCO ₃ , is the amount of HCO ₃ that must be added (mol/kg). The second parameter, H ⁺ , is the quantity of H ⁺ that must be added (mol/kg). The acid must be fortified with NaCl in order to have the same salinity than seawater. <i>Addition of CO₃ and acid</i> : carbonate, CO ₃ , and a strong acid are added. The first parameter, HCO ₃ , is the quantity of CO ₃ that must be added (mol/kg). The second parameter, H ⁺ , is the quantity of H ⁺ that must be added (mol/kg).
summary	Table summarizing the carbonate chemistry before and after applying each perturbation: pCO ₂ bubbling, mixing with high-CO ₂ seawater, addition of strong acid, and addition of bicarbonate/carbonate and strong acid.

Warnings

- It is recommended to use concentrated solutions of acid or base in order to add small volumes.
- The addition of strong acid does not simulate well natural ocean acidification (higher concentration of dissolved inorganic carbon at constant total alkalinity) since it generates a decrease in total alkalinity while dissolved inorganic carbon is kept constant.

- Other important advice is provided in Gattuso and Lavigne (2009), Schulz et al. (2009) and in the “Guide for Best Practices on Ocean Acidification Research and Data Reporting” (<https://www.iaea.org/sites/default/files/18/06/oa-guide-to-best-practices.pdf>)

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Author(s)

Heloise Lavigne and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Gattuso J.-P. and Lavigne H., 2009 Technical note: approaches and software tools to investigate the impact of ocean acidification. *Biogeosciences* **21**, 6:2121-2133.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993 The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Schulz K. G., Barcelos e Ramos J., Zeebe R. E. and Riebesell U., 2009 CO₂ perturbation experiments: similarities and differences between dissolved inorganic carbon and total alkalinity manipulations. *Biogeosciences* **6**, 2145-2153.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* 165, 66-67.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

See Also

[carb](#), [pgas](#), [pmix](#), [ppH](#), [pTA](#).

Examples

```
oa(flag=24, var1=384, var2=2325e-6, pCO2s=1e6, pCO2f=793, S=34.3, T=16,
P=0, pHscale="T", kf="pf", k1k2="1", ks="d", plot=TRUE, b="u74")
```

Om

Carbonate saturation state for magnesian calcites

Description

Calculates the calcium carbonate saturation state for magnesian calcite

Usage

```
Om(x, flag, var1, var2, k1k2='x', kf='x', ks="d", pHscale="T", b="u74")
```

Arguments

x	mole fraction of magnesium ions, note that the function is only valid for x ranging between 0 and 0.25
flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given flag = 6 pH and HCO ₃ given flag = 7 pH and CO ₃ given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO ₃ and CO ₃ given

	flag = 11 HCO3 and ALK given
	flag = 12 HCO3 and DIC given
	flag = 13 CO3 and ALK given
	flag = 14 CO3 and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "s20" from Sulpis et al. (2020), "sb21" from Shockman & Byrne (2021), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"

Details

It is important to note that this function is **only valid** for:

- Salinity = 35
- Temperature = 25 degrees Celsius
- Hydrostatic pressure = 0 bar (surface)
- Concentration of total phosphate = 0 mol/kg
- Concentration of total silicate = 0 mol/kg

Note that the stoichiometric solubility products with respect to Mg-calcite minerals have not been determined experimentally. The saturation state with respect to Mg-calcite minerals is therefore calculated based on ion activities, i.e.,

$$\Omega_x = \frac{\{Ca^{2+}\}^{1-x} \{Mg^{2+}\}^x \{CO_3\}^{2-}}{K_x}$$

The ion activity $\{a\}$ is calculated based on the observed ion concentrations $[C]$ multiplied by the total ion activity coefficient, γ_T , which has been determined experimentally or from theory (e.g. Millero & Pierrot 1998): $\{a\} = \gamma_T [C]$. Because a true equilibrium cannot be achieved with respect to Mg-calcite minerals, K_x represents a metastable equilibrium state obtained from what has been referred to as stoichiometric saturation (Thorstenson & Plummer 1977; a term not equivalent to the definition of the stoichiometric solubility product, see for example Morse et al. (2006) and references therein). In the present calculation calcium and magnesium concentrations were calculated based on salinity. Total ion activity coefficients with respect to Ca^{2+} , Mg^{2+} , and CO_3^{2-} were adopted from Millero & Pierrot (1998).

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al. (2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks_i, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_f, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

The function returns a list with

OmegaMgCa_biogenic

Mg-calcite saturation state for minimally prepared biogenic Mg-calcite.

OmegaMgCa_biogenic_cleaned

Mg-calcite saturation state for cleaned and annealed biogenic Mg-calcite.

Author(s)

Heloise Lavigne, Andreas J. Andersson and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Only the references related to the saturation state of magnesian calcite are listed below; the other references are listed under the carb function.

Andersson A. J., Mackenzie F. T., Nicholas R. B., 2008, Life on the margin: implications of ocean acidification on Mg-calcite, high latitude and cold-water marine calcifiers. *Marine Ecology Progress Series* **373**, 265-273.

Bischoff W. D., Mackenzie F. T. and Bishop F. C., 1987. Stabilities of synthetic magnesian calcites in aqueous solution: comparison with biogenic materials. *Geochimica et Cosmochimica Acta* **51**:1413-1423.

Millero F. J. and Pierrot D., 1998. A chemical equilibrium model for natural waters. *Aquatic Geochemistry* **4**, 153-199.

Morse J. W., Andersson A. J. and Mackenzie F. T., 2006. Initial responses of carbonate-rich shelf sediments to rising atmospheric pCO₂ and ocean acidification: Role of high Mg-calcites. *Geochimica et Cosmochimica Acta* **70**, 5814-5830.

Plummer L. N. and Mackenzie F. T., 1974. Predicting mineral solubility from rate data: application to the dissolution of magnesian calcites. *American Journal of Science* **274**:61-83.

Thorstenson D.C. and Plummer L.N., 1977. Equilibrium criteria for two component solids reacting with fixed composition in an aqueous phase-example: the magnesian calcites. *American Journal of Science* **277**, 1203-1233.

Examples

```
0m(x=seq(0.01, 0.252, 0.01), flag=8, var1=8.2, var2=0.00234,  
k1k2='x', kf='x', ks="d", pHscale="T", b="u74")
```

p2d

Converts pressure in dbar to depth in meters

Description

Converts pressure in dbar to depth in meters

Usage

```
p2d(pressure, lat=40)
```

Arguments

pressure	Pressure in dbar
lat	Latitude in degrees, N and S is irrelevant, default is 40o

Value

depth	Depth corresponding to the pressure given, in meters
-------	--

Author(s)

Heloise Lavigne and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Saunders P. M., 1981. Practical conversion of pressure to depth. *J. Phys. Oceanogr.* **11**: 573-574.

See Also

[d2p](#)

Examples

```
p2d(pressure=7686, lat=30)
```

*p2fCO2**Converts pCO2 (partial pressure in CO2) into fCO2 (fugacity of CO2)*

Description

Converts pCO2 (partial pressure in CO2) into fCO2 (fugacity of CO2)

Usage

`p2fCO2(T = 25, Patm=1, P=0, pCO2)`

Arguments

<code>T</code>	Temperature in degrees Celsius, default is 25oC
<code>Patm</code>	Surface atmospheric pressure in atm, default is 1 atm
<code>P</code>	Hydrostatic pressure in bar, default is 0 bar (surface)
<code>pCO2</code>	Partial pressure in CO2 in μ atm, the same units as that for the fugacity output

Value

`fCO2` Fugacity of CO2 in μ atm, the same units as that for the pCO2 input.

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Heloise Lavigne, Jean-Pierre Gattuso, and James Orr <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

See Also

[f2pCO2](#).

Examples

```
p2fCO2(T=25, Patm=0.97, P=0, pCO2=380)
```

p2xCO2

Converts partial pressure of CO2 to mole fraction of CO2

Description

Converts pCO2 (partial pressure of CO2) into xCO2 (mole fraction of CO2)

Usage

```
p2xCO2(S=35, T=25, Patm=1, pCO2)
```

Arguments

S	Salinity on the practical salinity scale, default is 35
T	Temperature in degrees Celsius, default is 25oC
Patm	Atmospheric pressure in atmospheres, default is 1.0 (this is not the hydrostatic pressure)
pCO2	Partial pressure of CO2 in μ atm

Details

The xCO2 (ppm) is computed from pCO2 (μ atm) using the following equation: $xCO2 = pCO2 / (Patm - pH2O)$, where pH2O is the vapor pressure of seawater computed following best practices (Dickson et al., 2007). That computed pH2O is identical, when rounded to the 4th decimal place, with that computed by the equation from Weiss and Price (1980).

Value

xCO2	Mole fraction of CO2 in ppm.
------	------------------------------

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

James Orr <james.orr@lsce.ipsl.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R. (2007) Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Weiss, R. F. (1974) Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, **2**, 203-215.

Weiss, R. F. and Price, B. A. (1980) Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

See Also

[x2pCO2](#) and [vapress](#)

Examples

```
## Convert atmospheric pressure from mbar to atm
Patm_mbar = 1052                      # in millibar
Patm      = Patm_mbar / 1013.25 # in atm
## Compute xCO2 from pCO2
pCO2 = 380
xCO2 = p2xCO2(T=25, S=35, Patm=Patm, pCO2=pCO2)
print(xCO2)
## The result is 377.1546 ppm
```

Description

Calculates the changes in the saturation states of aragonite and calcite resulting from the manipulation of the calcium concentration

Usage

```
pCa(flag, var1, var2, Ca, S=35, T=20, P=0, Pt=0, Sit=0, k1k2="x",
kf="x", ks="d", pHscale="T", b="u74", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given
------	--

	flag = 6 pH and HCO ₃ given
	flag = 7 pH and CO ₃ given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO ₃ and CO ₃ given
	flag = 11 HCO ₃ and ALK given
	flag = 12 HCO ₃ and DIC given
	flag = 13 CO ₃ and ALK given
	flag = 14 CO ₃ and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO ₂ and pH given
	flag = 22 pCO ₂ and HCO ₃ given
	flag = 23 pCO ₂ and CO ₃ given
	flag = 24 pCO ₂ and ALK given
	flag = 25 pCO ₂ and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO ₂ in μatm
var2	Value of the second variable in mol/kg, except for pH
Ca	Calcium concentration in mol/kg
S	Salinity
T	Temperature in degrees Celsius
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg
Sit	Concentration of total silicate in mol/kg
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_f , the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

comment	The initial or final state water
S	Salinity
T	Temperature in degrees Celsius
P	Pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	pCO2, CO2 partial pressure (μ atm)
fCO2	fCO2, CO2 fugacity (μ atm)
HCO3	HCO3 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Ben-Yaakov S. and Goldhaber M. B., 1973 The influence of sea water composition on the apparent constants of the carbonate system. *Deep-Sea Research* **20**, 87-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl}(\text{s}) + 1/2\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Gattuso J.-P. and Lavigne H, 2009 Perturbation experiments to investigate the impact of ocean acidification: approaches and software tools. *Biogeosciences* **6**, 4413-4439.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* 165, 66-67.

Examples

```
pCa(flag=15, var1=2302e-6, var2=2050e-6, Ca=0.01028, S=35, T=20, P=0,
Pt=0, Sit=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74") # with normal Ca concentration
pCa(flag=15, var1=2302e-6, var2=2050e-6, Ca=0.01028/2, S=35, T=20, P=0,
Pt=0, Sit=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74") # with 0.5 * Ca concentration
```

<i>pCO2insi</i>	<i>pCO2 at in situ temperature</i>
-----------------	------------------------------------

Description

Correction to compensate for the difference in temperature between the temperature of measurement and in situ temperature.

Usage

```
pCO2insi(pCO2lab = 400, Tlab = 20, SST = 19)
```

Arguments

<i>pCO2lab</i>	Partial pressure of CO2 measured in the lab in μatm
<i>Tlab</i>	Temperature of measurement in the lab in in degrees Celsius, default is 20oC
<i>SST</i>	Temperature in degrees Celsius, default is 19oC

Value

<i>pCO2insi</i>	Partial pressure of CO2 at in situ temperature in μatm , the same units as that for the pCO2 input.
-----------------	--

Note

The empirical correction applied is from Takahashi (1993) as recommended by Pierrot et al. (2009)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Pierrot D., Neill C., Sullivan K., Castle R., Wanninkhof R., LÃ¼ger H., Johannessen T., Olsen A., Feely R. A. and Cosca C. E., 2009. Recommendations for autonomous underway pCO2 measuring systems and data-reduction routines. *Deep-Sea Res. II* **56**, 512-522.

Takahashi T., Olafsson J., Goddard J. G., Chipman D. W. and Sutherland S. C., 1993. Seasonal variation of CO2 and nutrients in the high-latitude surface oceans: a comparative study. *Glob. Biogeochem. Cycles* **7**, 843-878.

See Also

[f2pCO2](#), [fCO2insi](#).

Examples

```
pCO2insi(pCO2lab = 400, SST = 15, Tlab = 16)
```

Pcoeffs*Coefficients used for pressure-correcting the equilibrium constants*

Description

Pressure corrections are based on the following equations:

$$\ln \frac{K_i^P}{K_i^0} = -\frac{\Delta V_i}{RT} \cdot P + 0.5 \frac{\Delta K_i}{RT} \cdot P^2$$

with

$$\Delta V_i = a_0 + a_1 T + a_2 T^2$$

and

$$\Delta K_i = b_0 + b_1 T + b_2 T^2$$

The variables are:

- K indicating the type of equilibrium constant
- coefficient a_0
- coefficient a_1
- coefficient a_2
- coefficient b_0
- coefficient b_1
- coefficient b_2

Usage**Pcoeffs****Format**

A data frame with 15 rows and 7 variables

Details

For Kb, to be consistent with Millero (1979) a2 was changed to -2.608e-3 instead of 2.608e-3 (value given in Millero, 1995) For Kw, coefficients are from Millero (1983).

Source

Millero F. J., 1979 The thermodynamics of the carbonate system in seawater. *Geochemica et Cosmochimica Acta* **43**: 1651-1661.

Millero F. J., 1983 Influence of pressure on chemical processes in the sea. pp. 1-88. In J. P. Riley and R. Chester (eds.), *Chemical Oceanography*. Academic Press, New York.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

See Also

Pcorrect

Pcorrect*Pressure correction of equilibrium constants*

Description

Computes the pressure correction of the equilibrium constants

Usage

```
Pcorrect(Kvalue, Ktype, T=25, S=35, P=0, pHscale="T",
          kconv2ScaleP0="x", kconv2Scale="x", warn="y")
```

Arguments

Kvalue Value of the constant at P=0 (hydrostatic pressure in bar, surface = 0)

Ktype Name of the constant,

- K1 First dissociation constant of carbonic acid (mol/kg)
- K2 Second dissociation constant of carbonic acid (mol/kg)
- Kb Dissociation constant of boric acid (mol/kg)
- Kw Ion product of water (mol²/kg²)
- Ks Stability constant of hydrogen sulfate (mol/kg)
- Kf Stability constant of hydrogen fluoride (mol/kg)
- Kspc Solubility product of calcite (mol/kg)
- Kspa Solubility product of aragonite (mol/kg)
- K1p First dissociation constant of phosphoric acid (mol/kg)
- K2p Second dissociation constant of phosphoric acid (mol/kg)
- K3p Third dissociation constant of phosphoric acid (mol/kg)
- Khs Dissociation constant of hydrogen sulfide(mol/kg)
- Kn Dissociation constant of ammonium (mol/kg)
- Ks1 Dissociation constant of Si(OH)₄ (mol/kg)
- K2s1 Second dissociation constant of Si(OH)₄ (mol/kg)

T Temperature in degrees Celsius, default is 25oC

S Salinity, default is 35

P Hydrostatic pressure in bar (surface = 0), default is 0

pHscale pHscale of the constant given in Kvalue

kconv2ScaleP0 Conversion factor from the pH scale selected to the SWS (or free for Kf) scale at pressure zero. It is advised to use default value "x", in which case it is computed when required. (may slow down the computation)

kconv2Scale	Conversion factor from the pH scale selected to the SWS (or free for Kf) scale at the hydrostatic pressure value indicated. It is advised to use default value "x", in which case it is computed when required. (may slow down the computation)
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".

Details

- The pressure correction is applied on the seawater scale for K1, K2, K1p, K2p, K3p, Kb, Khs, Kn, Ks1, K2s1 and Kw. Hence the K value is first converted on the seawater scale if needed. After pressure correction, the constant is converted back to the initial pH scale.
- The pressure correction is applied on the free scale for Kf.
- There is no issue of pH scale for Ks, Kspa and Kspc.

Value

The equilibrium constant given in argument but after pressure correction

Author(s)

Heloise Lavigne and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59** 661-677.

See Also

Pcoeffs

Examples

```
k10 <- K1(T=25, P=0, S=35)
Pcorrect(Kvalue=k10, Ktype="K1", P=300, T=25, S=35, pHscale="T")
```

Description

Calculates the carbonate chemistry after changes in pCO₂ generated by gas bubbling

Usage

```
pgas(flag, var1, var2, pCO2g, S=35, T=20, P=0, Pt=0, Sit=0, k1k2="x",
kf="x", ks="d", pHscale="T", b="u74", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
pCO2g	CO2 partial pressure of the gas used for bubbling in μatm
S	Salinity
T	Temperature in degrees Celsius
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg
Sit	Concentration of total silicate in mol/kg
k1k2	"1" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "1", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".

ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.

- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks_i, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_f, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

comment	The initial or final state water
S	Salinity
T	Temperature in degrees Celsius
P	Pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	pCO2, CO2 partial pressure (μ atm)
fCO2	fCO2, CO2 fugacity (μ atm)
HCO3	HCO3 concentration (mol/kg)

CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Jean-Pierre Gattuso and Heloise Lavigne <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl}(s) + 1/2\text{H}_2(g) = \text{Ag}(s) + \text{HCl}(aq)$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Gattuso J.-P. and Lavigne H, 2009 Perturbation experiments to investigate the impact of ocean acidification: approaches and software tools. *Biogeosciences* **6**, 4413-4439.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica et Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

Examples

```
pgas(flag=15, var1=2302e-6, var2=2050e-6, pCO2g=750, S=35, T=20, P=0,  
Pt=0, Sit=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
```

pH

Potentiometric pH

Description

Calculation of potentiometric pH

Usage

```
pH(Ex=-67,Etris=-72.4,S=35,T=25)
```

Arguments

Ex	e.m.f. of the seawater sample in mV, default is 67
Etris	e.m.f. of the TRIS buffer in mV, default is -72.4
S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC

Details

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

pH Potentiometric pH (in mol/kg on the total scale)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

See Also

[tris](#), [amp](#), [pHslope](#).

Examples

```
##Example from Dickson et al. (2007)
pH(Ex=-67,Etris=-72.4,S=35,T=25)
```

pHconv*Conversion of pH*

Description

Converts pH from one scale to another one chosen between the total scale, the free scale and the seawater scale

Usage

```
pHconv(flag=1, pH=8.10, S=35, T=25, P=0, ks="d")
```

Arguments

flag	choice of the type of conversion : flag=1: seawater scale to total scale flag=2: free scale to total scale flag=3: total scale to seawater scale flag=4: total scale to free scale flag=5: seawater scale to free scale flag=6: free scale to seawater scale default is flag=1
pH	Enter the value of pH which need to be converted, default is 8.100
S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0
ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"

Details

The Dickson (1990) constant is recommended by the Guide to Best Practices for Ocean CO₂ Measurements (2007). It is critical to consider that each formulation is valid in specific ranges of temperature and salinity:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Value

The function returns the values of pH converted

Author(s)

Heloise Lavigne and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A.G. and F.J. Millero, 1987 A comparison of the equilibrium constants for the dissociation of carbonic acid in seawater media. *Deep-Sea Research* **34**:1733-1743.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl}(\text{s}) + 1/2\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

See Also

[kconv](#).

Examples

```
##To convert pH=8.10 from the seawater scale to the total scale
##at salinity=35, temperature=25°C and atmospheric pressure:
pHc <- pHconv(flag=1, pH=8.10, S=35, T=25, P=0, ks="d")

##note that pHc is the value of the pH converted in total scale

## By using vectors
## to convert the pH values : 8, 8.05, 8.10, 8.15, 8.20
## from the free to the total scale

pH <- c(8, 8.05, 8.10, 8.15, 8.20)
pHc <- pHconv(flag=2, pH=pH, S=35, T=25, P=0, ks="d")

## note that pHc is a vector containing the value of the pH converted
## to the total scale
```

pHinsi

pH at in situ temperature and pressure

Description

pH at in situ temperature and pressure

Usage

```
pHinsi(pH=8.2, ALK=2.4e-3, Tinsi=20, Tlab=25, Pinsi=0, S=35, Pt=0, Sit=0,
      k1k2 = "x", kf = "x", ks="d", pHscale = "T", b="u74", eos = "eos80",
      long = 1e+20, lat = 1e+20)
```

Arguments

pH	pH measured in the laboratory
ALK	ALK, total alkalinity (mol/kg)
Tinsi	In situ temperature in degrees Celsius
Tlab	Measurement temperature in degrees Celsius
Pinsi	In situ hydrostatic pressure in bar (surface = 0)
S	Salinity
Pt	value of the concentration of total phosphate in mol/kg
Sit	the value of the total silicate in mol/kg
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	"l10" for computing boron total from the Lee et al. (2010) formulation or "u74" for using the Uppstrom (1974) formulation, default is "u74"
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al. (2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

pH pH at in situ temperature and pressure

Author(s)

Jean-Pierre Gattuso, <gattuso@obs-vlfr.fr>

References

Hunter K. A., 1998. The temperature dependence of pH in surface seawater. *Deep-Sea Research (Part I, Oceanographic Research Papers)* **45**(11):1919-1930.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

Examples

`pHinsi(pH=8.2,ALK=2.4e-3,Tinsi=25,Tlab=25,Pinsi=200,S=35,Pt=0,Sit=0)`

pHslope	<i>Slope of the calibration curve of a pH electrode</i>
---------	---

Description

Slope of the calibration curve of a pH electrode (percent of theoretical slope)

Usage

```
pHslope(Etris=-72.4, Eamp=4.9, S=35, T=25)
```

Arguments

Etris	e.m.f. of the TRIS buffer in mV, default is -72.4
Eamp	e.m.f. of the AMP buffer in mV, default is 4.9
S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC

Details

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

pHslope Slope of the calibration curve (in percent of theoretical slope)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication 3*, 1-191.

See Also

[tris](#), [amp](#), [pH](#).

Examples

```
##Example from Dickson et al. (2007)
pHslope(Etris=-72.4, Eamp=4.9, S=35, T=25)
```

pHspec*Calculates pHT from results of spectrophotometric measurements*

Description

Calculates pH of a water sample from absorbance ratios R, obtained from spectrophotometric measurements with pH indicator dyes (on the total scale in mol/kg-soln)

Usage

```
pHspec(S=35, T=25, R=1, d="mCP", k="m18", warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
R	Absorbance ratio, default is 1
d	Dye used for spectrophotometric measurement, default is "mCP"
k	"m18" for using mCP characterization by Mueller and Rehder (2018)
warn	"y" to show warnings when S and/or T go beyond the valid range for the chosen d and k; "n" to suppress warnings. The default is "y".

Details

The model used to calculate the return value of this function is based on experimental data. It is critical to consider that the formulation refers to the conditions studied during the characterization experiment and is only valid for the studied ranges of temperature and salinity:

- Mueller and Rehder (2018): S ranging between 0 and 40, T ranging between 5 and 35oC, and the dye used being m-Cresol purple (mCP) with R referring to the ratio of absorbances at wavelengths 578 and 434 nm.

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

pHspec	The function returns the pH value of a water sample from absorbance ratios R, obtained from spectrophotometric measurements with pH indicator dyes (on the total scale in mol/kg-soln)
--------	--

Author(s)

Jens Daniel Mueller <jens.mueller@io-warnemuende.de> Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Mueller, J. D. and Rehder, G., 2018 Metrology for pH Measurements in Brackish Waters - Part 2: Experimental Characterization of Purified meta-Cresol Purple for Spectrophotometric pHT Measurements. *Frontiers in Marine Science* **5**:177, 1-9. <https://doi.org/10.3389/fmars.2018.00177>

See Also

[amp](#), [pHslope](#), [pH_tris](#),

Examples

```
##Example should give test value pHT = 7.6713
pHspec(S=35, T=25, R=1, d="mCP", k="m18", warn="y")
```

pmix

pmix

Description

Calculates the carbonate chemistry after mixing of two water samples with different pCO2

Usage

```
pmix(flag, var1, var2, pCO2s, wf, S=35, T=20, P=0, Pt=0, Sit=0, k1k2="x",
kf="x", ks="d", pHscale="T", b="u74", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given
------	---

	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	Value of the first variable in mol/kg except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg except for pH
pCO2s	Partial pressure of the high CO2 component in μatm
wf	Weight fraction of the high CO2 seawater per kg seawater
S	Salinity
T	Temperature in degrees Celsius
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg
Sit	Concentration of total silicate in mol/kg
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickon (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al. (2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

comment	The initial or final state water
S	Salinity
T	Temperature in degrees Celsius
P	Pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	pCO2, CO2 partial pressure (μatm)
fCO2	fCO2, CO2 fugacity (μatm)
HCO3	HCO3 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Jean-Pierre Gattuso and Heloise Lavigne<gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Gattuso J.-P. and Lavigne H., 2009. Perturbation experiments to investigate the impact of ocean acidification: approaches and software tools. *Biogeosciences* **6**, 4413-4439.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

Examples

```
pmix(flag=24, var1=384, var2=2302e-6, pCO2s=1e6, wf=0.003, S=34.3,
T=16, P=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
```

ppH

ppH

Description

Calculates the carbonate chemistry after pH manipulations through addition of acid or base

Usage

```
ppH(flag, sys, var1, var2, pCO2a, vol, N, S=35, T=20, P=0, Pt=0,
  Sิต=0, pHscale="T", k1k2="x", kf="x", ks="d", eos = "eos80",
  long = 1e+20, lat = 1e+20)
```

Arguments

flag	Select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
sys	0 if the manipulation is carried out in a system closed to the atmosphere or 1 if its is carried out in a system open to the atmosphere
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μ atm
var2	Value of the second variable in mol/kg, except for pH
pCO2a	CO2 partial pressure in the atmosphere pCO2 in μ atm. It is only used in systems open to the atmosphere (i.e. when sys=1)
vol	Volume of acid or base added in liter. By convention, it is given a negative sign for acid additions and a positive sign for base additions. The acid must be fortified with NaCl in order to have the same salinity than seawater.
N	Normality of the acid or base in mol/kg
S	Salinity
T	Temperature in degrees Celsius
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg

Sit	Concentration of total silicate in mol/kg
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	choice of pH scale: "T" for using the total scale, "F" for using the free scale and "SWS" for using the seawater scale, default is total scale
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.

- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

comment	The initial or final state water
S	Salinity
T	Temperature in degrees Celsius
P	Pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	pCO2, CO2 partial pressure (μ atm)
fCO2	fCO2, CO2 fugacity (μ atm)
HCO3	HCO3 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Jean-Pierre Gattuso and Heloise Lavigne<gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Gattuso J.-P. and Lavigne H., 2009 Perturbation experiments to investigate the impact of ocean acidification: approaches and software tools. *Biogeosciences* **6**, 4413-4439.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHE: 149 (2013) 8-22], Marine Chemistry 165, 66-67.

See Also

[buffer](#).

Examples

```
ppH(flag=24, sys=0, var1=384, var2=2302e-6, pCO2a=384, vol=-12e-3,
N=0.01, S=34.3, T=16, P=0, pHscale="T", kf="pf", k1k2="1", ks="d")
```

```
ppH(flag=24, sys=1, var1=384, var2=2302e-6, pCO2a=384, vol=-12e-3,
N=0.01, S=34.3, T=16, P=0, pHscale="T", kf="pf", k1k2="1", ks="d")
```

psi

Molar ratio of CO₂ released vs CaCO₃ precipitated

Description

Returns the molar ratio of CO₂ released vs CaCO₃ precipitated described by Frankignoulle et al. (1994).

Usage

```
psi(flag, var1, var2, S=35, T=20, Patm, P=0, Pt=0, Sit=0, pHscale="T",
kf="x", k1k2="x", ks="d", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given flag = 6 pH and HCO ₃ given flag = 7 pH and CO ₃ given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO ₃ and CO ₃ given flag = 11 HCO ₃ and ALK given flag = 12 HCO ₃ and DIC given flag = 13 CO ₃ and ALK given
------	--

	flag = 14 CO3 and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given
var1	enter value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	enter value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
pHscale	choice of pH scale: "T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al. (2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

psi ratio of CO₂ released vs CaCO₃ precipitated (mol/mol)

Author(s)

Jean-Pierre Gattuso and Heloise Lavigne <gattuso@obs-vlfr.fr>

References

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication 3*, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Frankignoulle M., 1994 A complete set of buffer factors for acid/base CO₂ system in seawater. *Journal of Marine Systems* **5**, 111-118.

Frankignoulle M., Canon C. and Gattuso J.P., 1994 Marine calcification as a source of carbon dioxide- Positive feedback of increasing atmospheric CO₂. *Limnology and Oceanography* **2**, 458-462.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to "The free proton concentration scale for seawater pH", [MARCHE: 149 (2013) 8-22], Marine Chemistry 165, 66-67.

See Also

[speciation](#).

Examples

```
## Calculation using the numerical example given in Frankignoulle et al. (1994)
psi(flag=24, var1=350, var2=2400e-6, S=35, T=25, P=0, Pt=0,
Sit=0, pHscale="T", kf="pf", k1k2="l", ks="d")
```

pTA

pTA

Description

Calculates the carbonate chemistry following addition of CO_3^{2-} or HCO_3^-

Usage

```
pTA(flag, sys=0, var1, var2, pCO2a, co3, hco3, S=35, T=20, P=0,
Pt=0, Sit=0, k1k2="x", kf="x", ks="d", pHscale="T", b="u74",
eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO2 given flag = 2 CO2 and HCO3 given flag = 3 CO2 and CO3 given flag = 4 CO2 and ALK given flag = 5 CO2 and DIC given flag = 6 pH and HCO3 given flag = 7 pH and CO3 given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO3 and CO3 given flag = 11 HCO3 and ALK given flag = 12 HCO3 and DIC given flag = 13 CO3 and ALK given flag = 14 CO3 and DIC given flag = 15 ALK and DIC given flag = 21 pCO2 and pH given flag = 22 pCO2 and HCO3 given flag = 23 pCO2 and CO3 given flag = 24 pCO2 and ALK given flag = 25 pCO2 and DIC given
------	--

sys	0 if the manipulation is carried out in a system closed to the atmosphere or 1 if its is carried out in a system open to the atmosphere
var1	Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg, except for pH
pCO2a	CO2 partial pressure in the atmosphere pCO2 in μatm . It is only used in systems open to the atmosphere (i.e. when sys=1)
co3	Amount of CO_3^{2-} added in mol kg^{-1}
hco3	Amount of HCO_3^{2-} added in mol kg^{-1}
S	Salinity
T	Temperature in degrees Celsius
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg
Sit	Concentration of total silicate in mol/kg
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990), "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al. (2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For K0:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

comment	The initial or final state water
S	Salinity
T	Temperature in degrees Celsius
P	Pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	pCO2, CO2 partial pressure (μatm)
fCO2	fCO2, CO2 fugacity (μatm)
HCO3	HCO3 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl(s)} + 1/2\text{H}_2\text{g} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Gattuso J.-P. and Lavigne H., 2009 Perturbation experiments to investigate the impact of ocean acidification: approaches and software tools. *Biogeosciences* **6**, 4413-4439.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Millero F. J., 1995 Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010 Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006 Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHÉ: 149 (2013) 8-22], *Marine Chemistry* 165, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Examples

```
pTA(flag=24, sys=0, var1=384, var2=2302e-6, pCO2a=384, co3=260e-6,
hco3=1000e-6, S=34.3, T=16, P=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
```

```
pTA(flag=24, sys=1, var1=384, var2=2302e-6, pCO2a=384, co3=260e-6,
hco3=1000e-6, S=34.3, T=16, P=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
```

rho	<i>Density of seawater (kg/m3)</i>
-----	------------------------------------

Description

Calculates the density of seawater ($kg\ m^{-3}$)

Usage

```
rho(S = 35, T = 25, P = 0)
```

Arguments

S	Practical Salinity (PSS-78), default is 35
T	Temperature in degrees Celsius (ITS-90), default is 25oC
P	Hydrostatic pressure in bar (surface = 0), default is 0

Value

rho Density of seawater (kg/m3)

Author(s)

Aurelien Proye and Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Millero F. J. and Poisson A., 1981 International one-atmosphere equation of state of seawater. *Deep-Sea Research* **28A**, 625-629.

Examples

```
rho(35,25,0)
```

sa2sp_chem	<i>From absolute to practical salinity</i>
------------	--

Description

Converts from absolute to practical salinity (SP). Salinity conversion depends on total alkalinity as well as the concentrations of dissolved inorganic carbon, nitrate and silicate.

Usage

```
sa2sp_chem(SA, TA=2300e-6, DIC=2000e-6, NO3=0, SiOH4=0)
```

Arguments

SA	Absolute salinity in g/kg
TA	Total alkalinity, in mol/kg, default is 2300 μ mol/kg
DIC	Dissolved inorganic carbon concentration in mol/kg, default is 2000 μ mol/kg
NO3	Total nitrate concentration in mol/kg, default is 0
SIOH4	Total silicate concentration in mol/kg, default is 0

Details

Convert from absolute to practical salinity from carbon system parameters and ion concentration which most affect water density anomalies.

Value

SP	Practical salinity (in psu)
----	-----------------------------

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., Wright D. G. and Millero F. J., 2011. The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* **7**, 363-387.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

sp2sa_chem does the reverse, sa2sp_geo

Examples

```
# Calculate the practical salinity of a sample with Absolute Salinity of 35 g/kg,  
# Total Alkalinity of 0.00234 mol/kg and DIC of 0.00202 mol/kg  
SP <- sa2sp_chem(SA=35, TA=0.00234, DIC=0.00202)
```

Description

Converts from absolute to practical salinity based on depth and geographic location.

Usage

```
sa2sp_geo(SA, P=0, long=1.e20, lat=1.e20)
```

Arguments

SA	Absolute salinity in g/kg
P	Sea water pressure in dbar
long	Longitude in decimal degrees [0 ... +360] or [-180 ... +180]
lat	Latitude in decimal degrees [-90 ... 90]

Details

This function is almost an alias of subroutine gsw_SP_from_SA from gsw package on which it relies. The only difference is in that depth and location are optional. If location is not given, or incomplete (either longitude or latitude missing), an arbitrary location is chosen: the mid equatorial atlantic ocean. Note that this implies an error on computed SA ranging from 0 up to 0.02 g/kg.

Value

SP	Practical salinity (psu)
----	--------------------------

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

sp2sa_geo does the reverse, sa2sp_chem

Examples

```
# Calculate the practical salinity of a sample whose absolute Salinity is 35,
# depth is 10 dbar and location is 188 degrees East and 4 degrees North.
SP <- sa2sp_geo(35, 10, 188, 4)
```

seacarb_test_P0	<i>Test data file (at P=0) to test the use of the carb function</i>
-----------------	---

Description

The variables are:

- Flag indicating which couple of variables is used
- Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
- Value of the second variable in mol/kg, except for pH
- Salinity
- Temperature in degrees Celsius
- Hydrostatic pressure in bar (surface = 0)
- Value of the concentration of total phosphate in mol/kg
- Value of the total silicate in mol/kg

Usage

seacarb_test_P0

Format

A data frame with 20 rows and 8 variables

Source

None, these data were invented for this purpose. The input variables were chosen in order to check that the carbonate chemistry is identical for all flags.

seacarb_test_P300 *Test data file (at P=300) to test the use of the carb function*

Description

The variables are:

- Flag indicating which couple of variables is used
- Value of the first variable in mol/kg, except for pH and for pCO2 in μatm
- Value of the second variable in mol/kg, except for pH
- Salinity
- Temperature in degrees Celsius
- Hydrostatic pressure in bar (P=300)
- Value of the concentration of total phosphate in mol/kg
- Value of the total silicate in mol/kg

Usage

seacarb_test_P300

Format

A data frame with 20 rows and 8 variables

Source

None, these data were invented for this purpose. The input variables were chosen in order to check that the carbonate chemistry is identical for all flags.

seaFET *Test seaFET data file*

Description

Short test file for using with functions sf_calc and sf_calib. It is an excerpt of a file produced by a SeaFET pH sensor.

Usage

seaFET

Format

A data frame with 8 variables (datetime, Eint, Eext, Salinity, Temperature, pHspectro, E0int25, E0ext25) and 10 rows

sf_calc*Calculation of calibrated pH for seaFET sensor*

Description

The function sf_calc() calculates pH time series (pHint_tot and pHext_tot) for SeaFET pH sensors, using calibration coefficients E0int25 and E0ext25 from the function sf_calib(). Both functions are R-adaptations from MATLAB scripts published by Bresnahan et al. (2014).

Usage

```
sf_calc(calEint=0.0865, calEext= -0.93, E0int25 =-0.39,
E0ext25=-1.46, calT=16.2, calSal=35.6)
```

Arguments

calEint	EINT (V), default is 0.0865
calEext	EEXT (V), default is -0.93
E0int25	Coeficient of calibration related to the internal sensor and obtained via sf_calib function, default is -0.39. If time-serie, we use the mean per periode of deployment
E0ext25	Coeficient of calibration related to the external sensor and obtained via sf_calib function, default is -1.46. If time-serie, we use the mean per periode of deployment
calT	Temperature in degrees Celsius, default is 16.2
calSal	Salinity, default is 35.6

Details

Input values should be vectors of equal length. E0int25 and E0ext25 should be constant throughout the time series. When multiple reference samples are available for one SeaFET deployment, mean E0int25 and mean E0ext25 should be calculated and used in sf_calc(). Each unique SeaFET deployment requires a new calculation of mean E0int25 and mean E0ext25 based on reference pH samples (total hydrogen ion scale). For detailed SeaFET calibration instructions and recommendations see Bresnahan et al. (2014) and Rivest et al. (2016).

Value

This function returns a dataframe comprising 2 variables:

pHint_tot	Calibrated pH of the internal sensor at in situ temperature.
pHext_tot	Calibrated pH of the external sensor at in situ temperature.

Author(s)

Samir Alliouane, Lydia Kapsenberg, Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Bresnahan, P. J. J., Martz, T. R., Takeshita, Y., Johnson, K. S., and LaShomb, M., 2014. Best practices for autonomous measurement of seawater pH with the Honeywell Durafet. *Methods Oceanogr.* **9**, 44-60.

Martz, T.R., Connery, J.G., and Johnson, K.S., 2010. Testing the Honeywell Durafet for seawater pH applications. *Limnol. Oceanogr. Meth.* **8**, 172-184.

Rivest, E.B., O'Brien, M., Kapsenberg, L., Gotschalk, C.C., Blanchette, C.A., Hoshijima, U., and Hofmann, G.E., 2016. Beyond the benchtop and the benthos: dataset management planning and design for time series of ocean carbonate chemistry associated with Durafet(c)-based pH sensors. *Ecological Informatics* **36**, 209-220.

See Also

[sf_calib](#).

Examples

```
sf_calc(calEint=0.0865, calEext= -0.93, E0int25 =-0.39,
E0ext25=-1.46, calT=16.2, calSal=35.6)

## Using the test file seaFET
sf_calc(calEint=seaFET$Eint, calEext=seaFET$Eext,
E0int25=seaFET$E0int25, E0ext25=seaFET$E0ext25,
calT=seaFET$Temperature, calSal=seaFET$Salinity)
```

sf_calib

Calibration coefficients for seaFET sensor

Description

Calibration coefficients E0INT,25, E0EXT,25

Usage

```
sf_calib(calEint=0.0865, calEext=-0.93, calpH=8.132, calT=16.2, calSal=35.6)
```

Arguments

calEint	EINT (V), default is 0.0865
calEext	EEXT (V), default is -0.93
calpH	spectrophotometric pH in Total scale, default is 8.132
calT	Temperature in degrees Celsius, default is 16.2
calSal	Salinity, default is 35.6

Details

Outputs E0INT25 and E0EXT25 must be calculated for each reference sample collected during a SeaFET deployment. Multiple E0INT25 and E0EXT25 may be calculated if there is more than one reference sample for a given deployment. As such, arguments can be given as unique numbers or as vectors (vectors should be of the same length).

It is critical that Eint (calEint) and Eext (calEext) recorded by the SeaFET match reference sample measurements of temperature (calT), salinity (calSal), and spectrophotometric pH (calpH, total hydrogen ion scale) taken at the same time. Note that SeaFET temperature measurements may require calibration via an applied offset. When possible, calibrated CTD temperature and salinity measurements may be used, while spectrophotometric pH measurements always require discrete 'reference' water samples (unless in situ, certified, seawater-based, Tris pH buffer is used). The accepted time offset between collection of reference samples and SeaFET measurements depends on the hydrology and pH variability of the location. For detailed SeaFET calibration instructions and recommendations see Bresnahan et al. (2014) and Rivest et al. (2016).

Value

This function returns a dataframe comprising 2 variables:

E0int25	Calibration coefficients of the internal sensor at 25oC.
E0ext25	Calibration coefficients of the external sensor at 25oC.

Author(s)

Samir Alliouane, Lydia Kapsenberg, Jean-Pierre Gattuso <gattuso@obs-vlfr.fr>

References

Bresnahan, P. J. J., Martz, T. R., Takeshita, Y., Johnson, K. S., and LaShomb, M., 2014. Best practices for autonomous measurement of seawater pH with the Honeywell Durafet. *Methods Oceanogr.* **9**, 44-60.

Martz, T.R., Connery, J.G., and Johnson, K.S., 2010. Testing the Honeywell Durafet for seawater pH applications. *Limnol. Oceanogr. Meth.* **8**, 172-184.

Rivest, E.B., O'Brien, M., Kapsenberg, L., Gotschalk, C.C., Blanchette, C.A., Hoshijima, U., and Hofmann, G.E., 2016. Beyond the benchtop and the benthos: dataset management planning and design for time series of ocean carbonate chemistry associated with Durafet(c)-based pH sensors. *Ecological Informatics* **36**, 209-220.

See Also

[sf_calc.](#)

Examples

```
sf_calib(calEint=0.0865, calEext=-0.93, calpH=8.132, calT=16.2, calSal=35.6)

## Using the test file seaFET
sf_calib(calEint=seaFET$Eint, calEext=seaFET$Eext,
```

```
calpH=seaFET$pHspectro, calT=seaFET$Temperature,
calSal=seaFET$Salinity)
```

sir

Parameters of the seawater carbonate system including the substrate-inhibitor-ratio (SIR)

Description

This function is a modification of *carb*. It returns parameters of the seawater carbonate system, as well as additional parameters relevant to the substrate-inhibitor-ratio (SIR).

Usage

```
sir(flag, var1, var2, S = 35, T = 25, Patm = 1, P = 0,
     Pt = 0, Sit = 0, k1k2 = "x", kf = "x", ks = "d", pHscale = "T",
     b = "u74", gas = "potential", warn = "y", eos = "eos80",
     long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given flag = 3 CO ₂ and CO ₃ given flag = 4 CO ₂ and ALK given flag = 5 CO ₂ and DIC given flag = 6 pH and HCO ₃ given flag = 7 pH and CO ₃ given flag = 8 pH and ALK given flag = 9 pH and DIC given flag = 10 HCO ₃ and CO ₃ given flag = 11 HCO ₃ and ALK given flag = 12 HCO ₃ and DIC given flag = 13 CO ₃ and ALK given flag = 14 CO ₃ and DIC given flag = 15 ALK and DIC given flag = 21 pCO ₂ and pH given flag = 22 pCO ₂ and HCO ₃ given flag = 23 pCO ₂ and CO ₃ given flag = 24 pCO ₂ and ALK given flag = 25 pCO ₂ and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO ₂ in μatm
var2	Value of the second variable in mol/kg, except for pH

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35oC and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33oC and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

Calculates the "substrate-inhibitor ratio (SIR)" (i.e. $[\text{HCO}_3^-]/[\text{H}^+]$) which is used as a metric to predict carbonate chemistry dependency of biotic calcium carbonate formation (Bach, 2015). Please note that hydrogen ion concentrations in the output $[\text{HCO}_3^-]/[\text{H}^+]$ are on the *free scale*, regardless

of the input pH scale. We have included pH and $[H^+]$ outputs on multiple scales as a teaching exercise to show how calculating the SIR on different scales changes its meaning.

The function is based on *carb* and therefore also returns parameters of the seawater carbonate system.

***carb* details:**

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.

- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μ atm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μ atm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μ atm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μ atm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)

fCO2insitu	"in situ" fCO ₂ , CO ₂ fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
HC03	HCO ₃ concentration (mol/kg)
CO3	CO ₃ concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state
SIR	substrate inhibitor ratio: [HCO ₃]/[H _{free}] (mol: μ mol)
H_free	H ion concentration on free scale (μ mol/kg)
H_sws	H ion concentration on seawater scale (μ mol/kg)
H_t	H ion concentration on total scale (μ mol/kg)
pH_free	pH calculated on the free scale
pH_sws	pH calculated on the seawater scale
pH_t	pH calculated on the total scale

Note

Warning: pCO₂ estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Kimberlee Baldry, Lennart Bach and Jean-Pierre Gattuso, <gattuso@obs-vlfr.fr>

References

Bach L. T., 2015 Reconsidering the role of carbonate ion concentration in calcification by marine organisms. *Biogeosciences* **12**, 4939-4951.

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: AgCl(s) + 1/2H₂(g) = Ag(s) + HCl(aq), and the standard acidity constant of the ion HSO₄ in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen ion concentration in seawater from 5 to 40°C: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21**, 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to “The free proton concentration scale for seawater pH”, [MARCHE: 149 (2013) 8-22], *Marine Chemistry* **165**, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```

## With a couple of variables
sir(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")

```

```

k1k2 <- c("1", "1", "1")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
sir(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P,
  Pt=Pt, Sit=Sit, kf=kf, k1k2=k1k2, pHscale=pHscale, b=b)

## Test with all flags
flag <- c((1:15), (21:25))
var1 <- c(8.200000, 7.308171e-06, 7.308171e-06, 7.308171e-06,
  8.2, 8.2, 8.2, 8.2, 0.001646857, 0.001646857, 0.001646857, 0.0002822957,
  0.0002822957, 0.00234, 258.2164, 258.2164, 258.2164, 258.2164 )
var2 <- c(7.308171e-06, 0.001646857, 0.0002822957, 0.00234, 0.001936461,
  0.001646857, 0.0002822957, 0.00234, 0.001936461, 0.0002822957,
  0.00234, 0.001936461, 0.00234, 0.001936461, 0.001936461, 8.2,
  0.001646857, 0.0002822957, 0.00234, 0.001936461)
sir(flag=flag, var1=var1, var2=var2)

## Test using a data frame
data(seacarb_test_P0) #test data set for P=0 (surface)
tab <- seacarb_test_P0[14:19,]

## method 1 using the column numbers
sir(flag=tab[[1]], var1=tab[[2]], var2=tab[[3]], S=tab[[4]], T=tab[[5]],
P=tab[[6]], Sit=tab[[8]], Pt=tab[[7]])

## method 2 using the column names
sir(flag=tab$flag, var1=tab$var1, var2=tab$var2, S=tab$S, T=tab$T,
P=tab$P, Sit=tab$Sit, Pt=tab$Pt)

```

sir_b

Parameters of the seawater carbonate system with boron addition, including the substrate-inhibitor-ratio (SIR)

Description

This function is a modification of *carbb*. It returns parameters of the seawater carbonate system when boron is added, as well as additional parameters relevant to the SIR.

Usage

```
sir_b(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
  k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential", badd=0,
  warn="y", eos = "eos80", long = 1e+20, lat = 1e+20)
```

Arguments

flag	select the couple of variables available. The flags which can be used are: flag = 1 pH and CO ₂ given flag = 2 CO ₂ and HCO ₃ given
------	--

	flag = 3 CO ₂ and CO ₃ given
	flag = 4 CO ₂ and ALK given
	flag = 5 CO ₂ and DIC given
	flag = 6 pH and HCO ₃ given
	flag = 7 pH and CO ₃ given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO ₃ and CO ₃ given
	flag = 11 HCO ₃ and ALK given
	flag = 12 HCO ₃ and DIC given
	flag = 13 CO ₃ and ALK given
	flag = 14 CO ₃ and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO ₂ and pH given
	flag = 22 pCO ₂ and HCO ₃ given
	flag = 23 pCO ₂ and CO ₃ given
	flag = 24 pCO ₂ and ALK given
	flag = 25 pCO ₂ and DIC given
var1	Value of the first variable in mol/kg, except for pH and for pCO ₂ in μatm
var2	Value of the second variable in mol/kg, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg; set to 0 if NA
Sit	Concentration of total silicate in mol/kg; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)

b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
badd	Amount of boron added in mol/kg.
warn	"y" to show warnings when T or S go beyond the valid range for constants; "n" to suppress warnings. The default is "y".
eos	"teos10" to specify T and S according to Thermodynamic Equation Of Seawater - 2010 (TEOS-10); "eos80" to specify T and S according to EOS-80.
long	longitude of data point, used when eos parameter is "teos10" as a conversion parameter from absolute to practical salinity.
lat	latitude of data point, used when eos parameter is "teos10".

Details

Calculates the "substrate-inhibitor ratio (SIR)" (i.e. $[\text{HCO}_3^-]/[\text{H}^+]$) which is used as a metric to predict carbonate chemistry dependency of biotic calcium carbonate formation (Bach, 2015). Please note that hydrogen ion concentrations in the output $[\text{HCO}_3^-]/[\text{H}^+]$ are on the *free scale*, regardless of the input pH scale. We have included pH and $[\text{H}^+]$ outputs on multiple scales as a teaching exercise to show how calculating the SIR on different scales changes its meaning.

The function is based on *carbb* and therefore also returns parameters of the seawater carbonate system when boron is added.

carbb details:

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35°C.
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45°C.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35°C.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50°C.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50°C. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25°C.
- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45°C.

- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35°C. For K1, formulation is that of Waters et al.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8°C.
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50°C. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33°C.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45°C.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45°C.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40°C.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs and Ks, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kf, the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides Kf on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, Kf was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Ks, the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For Kn, The pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

long and lat are used as conversion parameters from absolute to practical salinity: when seawater is not of standard composition, practical salinity alone is not sufficient to compute absolute salinity and vice-versa. One needs to know the density. When long and lat are given, density is inferred from WOA silicate concentration at given location. When they are not, an arbitrary geographic point is chosen: mid equatorial Atlantic. Note that this implies an error on computed salinity up to 0.02 g/kg.

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μatm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μatm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μatm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μatm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μatm)
fCO2insitu	"in situ" fCO2, CO2 fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μatm)
HC03	HC03 concentration (mol/kg)
CO3	CO3 concentration (mol/kg)
DIC	DIC concentration (mol/kg)
ALK	ALK, total alkalinity (mol/kg)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state
SIR	substrate inhibitor ratio: [HCO3]/[HCO3]/[H_free] (mol: μmol)
H_free	H ion concentration on free scale ($\mu\text{mol/kg}$)
H_sws	H ion concentration on seawater scale ($\mu\text{mol/kg}$)
H_t	H ion concentration on total scale ($\mu\text{mol/kg}$)
pH_free	pH calculated on the free scale
pH_sws	pH calculated on the seawater scale
pH_t	pH calculated on the total scale

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

Kimberlee Baldry, Lennart Bach and Jean-Pierre Gattuso, <gattuso@obs-vlfr.fr>

References

Bach L. T., 2015 Reconsidering the role of carbonate ion concentration in calcification by marine organisms. *Biogeosciences* **12**, 4939-4951.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen ion concentration in seawater from 5 to 40oC: standard potentials at salinities from 20 to 45. *Analytical Chemistry* **49**, 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO₂ calculated from dissolved inorganic carbon, alkalinity, and equations for K1 and K2: validation based on laboratory measurements of CO₂ in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45oC. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```
## With a couple of variables
sir_b(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0,
Pt=0, Sit=0, pHscale="T", kf="pf", k1k2="1", ks="d", b="u74", badd=0)
```

sir_full

Parameters of the seawater carbonate system including the substrate-inhibitor-ratio (SIR) - extension of carb

Description

This function is a modification of carb. It returns parameters of the seawater carbonate system, including the ammonium and sulfide acid-base systems, as well as full acid-base speciation and additional parameters relevant to the SIR.

Usage

```
sir_full(flag, var1, var2, S=35, T=25, Patm=1, P=0, Pt=0, Sit=0,
k1k2="x", kf="x", ks="d", pHscale="T", b="u74", gas="potential",
NH4t=0, HSt=0)
```

Arguments

flag	select the couple of variables available. The flags which can be used are:
	flag = 1 pH and CO2 given
	flag = 2 CO2 and HCO3 given
	flag = 3 CO2 and CO3 given
	flag = 4 CO2 and ALK given
	flag = 5 CO2 and DIC given
	flag = 6 pH and HCO3 given
	flag = 7 pH and CO3 given
	flag = 8 pH and ALK given
	flag = 9 pH and DIC given
	flag = 10 HCO3 and CO3 given
	flag = 11 HCO3 and ALK given
	flag = 12 HCO3 and DIC given
	flag = 13 CO3 and ALK given
	flag = 14 CO3 and DIC given
	flag = 15 ALK and DIC given
	flag = 21 pCO2 and pH given
	flag = 22 pCO2 and HCO3 given
	flag = 23 pCO2 and CO3 given
	flag = 24 pCO2 and ALK given
	flag = 25 pCO2 and DIC given

var1	Value of the first variable in mol/kg-soln, except for pH and for pCO2 in μatm
var2	Value of the second variable in mol/kg-soln, except for pH
S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm, default is 1 atm
P	Hydrostatic pressure in bar (surface = 0)
Pt	Concentration of total phosphate in mol/kg-soln; set to 0 if NA
Sit	Concentration of total silicate in mol/kg-soln; set to 0 if NA
k1k2	"l" for using K1 and K2 from Lueker et al. (2000), "m02" from Millero et al. (2002), "m06" from Millero et al. (2006), "m10" from Millero (2010), "mp2" from Mojica Prieto et al. (2002), "p18" from Papadimitriou et al. (2018), "r" from Roy et al. (1993), "sb21" from Shockman & Byrne (2021), "s20" from Sulpis et al. (2020), and "w14" from Waters et al. (2014). "x" is the default flag; the default value is then "l", except if T is outside the range 2 to 35°C and/or S is outside the range 19 to 43. In these cases, the default value is "w14".
kf	"pf" for using Kf from Perez and Fraga (1987) and "dg" for using Kf from Dickson and Riley (1979 in Dickson and Goyet, 1994). "x" is the default flag; the default value is then "pf", except if T is outside the range 9 to 33°C and/or S is outside the range 10 to 40. In these cases, the default is "dg".
ks	"d" for using Ks from Dickson (1990) and "k" for using Ks from Khoo et al. (1977), default is "d"
pHscale	"T" for the total scale, "F" for the free scale and "SWS" for using the seawater scale, default is "T" (total scale)
b	Concentration of total boron. "l10" for the Lee et al. (2010) formulation or "u74" for the Uppstrom (1974) formulation, default is "u74"
gas	used to indicate the convention for INPUT pCO2, i.e., when it is an input variable (flags 21 to 25): "insitu" indicates it is referenced to in situ pressure and in situ temperature; "potential" indicates it is referenced to 1 atm pressure and potential temperature; and "standard" indicates it is referenced to 1 atm pressure and in situ temperature. All three options should give identical results at surface pressure. This option is not used when pCO2 is not an input variable (flags 1 to 15). The default is "potential".
NH4t	Concentration of total ammonium in mol/kg-soln; set to 0 if NA
HSt	Concentration of total hydrogen sulfide in mol/kg-soln; set to 0 if NA

Details

Calculates the "substrate-inhibitor ratio (SIR)" (i.e. $[\text{HCO}_3^-]/[\text{H}^+]$) which is used as a metric to predict carbonate chemistry dependency of biotic calcium carbonate formation (Bach, 2015). Please note that hydrogen ion concentrations in the output $[\text{HCO}_3^-]/[\text{H}^+]$ are on the *free scale*, regardless of the input pH scale. We have included pH and $[\text{H}^+]$ outputs on multiple scales as a teaching exercise to show how calculating the SIR on different scales changes its meaning.

The function is based on *carbfull* and therefore also returns parameters of the seawater carbonate system, including the ammonium and sulfide acid- base systems, as well as full acid-base speciation.

***carbfull* details:**

The Lueker et al. (2000) constants for K1 and K2, the Perez and Fraga (1987) constant for Kf and the Dickson (1990) constant for Ks are recommended by Dickson et al. (2007). It is, however, critical to consider that each formulation is only valid for specific ranges of temperature and salinity:

For K1 and K2:

- Roy et al. (1993): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Lueker et al. (2000): S ranging between 19 and 43 and T ranging between 2 and 35oC.
- Mojica Prieto et al. (2002): S ranging from 5 to 42 and T ranging between 0 and 45oC.
- Millero et al. (2002): S ranging from 34 to 37 and T ranging between -1.6 and 35oC.
- Millero et al. (2006): S ranging between 0.1 and 50 and T ranging between 1 and 50oC.
- Millero (2010): S ranging between 1 and 50 and T ranging between 0 and 50oC. Millero (2010) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Waters et al.(2014): S ranging between 1 and 50 and T ranging between 0 and 50oC. Waters (2014) provides a K1 and K2 formulation for the seawater, total and free pH scales. Therefore, when this method is used and if P=0, K1 and K2 are computed with the formulation corresponding to the pH scale given in the flag "pHscale".
- Papadimitriou et al. (2018): S ranging from 33 to 100 and T ranging between -6 to 25oC.
- Sulpis et al. (2020): S ranging from 30.7 to 37.6 and T ranging between -1.7 to 31.8oC.
- Shockman & Byrne (2021): for K2, S ranging from 19.6 to 41 and T ranging between 15 to 35oC. For K1, formulation is that of Waters et al.

For Kf:

- Perez and Fraga (1987): S ranging between 10 and 40 and T ranging between 9 and 33oC.
- Dickson and Riley (1979 in Dickson and Goyet, 1994): S ranging between 0 and 45 and T ranging between 0 and 45oC.

For Ks:

- Dickson (1990): S ranging between 5 and 45 and T ranging between 0 and 45oC.
- Khoo et al. (1977): S ranging between 20 and 45 and T ranging between 5 and 40oC.

The arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It is recommended to use either vectors with the same dimension or one vector for one argument and numbers for the other arguments.

Pressure corrections and pH scale:

- For K0, the pressure correction term of Weiss (1974) is used.
- For K1, K2, pK1, pK2, pK3, Kw, Kb, Khs, Ks and K2si, the pressure correction was applied on the seawater scale. Hence, if needed, values were first transformed from the total scale to the seawater scale, the pressure correction applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

- For K_f , the pressure correction was applied on the free scale. The formulation of Dickson and Riley (1979 in Dickson and Goyet, 1994) provides K_f on the free scale but that of Perez and Fraga (1987) provides it on the total scale. Hence, in that case, K_f was first transformed from the total scale to the free scale. With both formulations, the pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_s , the pressure correction was applied on the free scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).
- For K_n , the pressure correction was applied on the seawater scale. The pressure correction was applied as described by Millero (1995), and the value was transformed back to the required scale (T, F or SWS).

Value

The function returns a data frame containing the following columns:

S	Salinity
T	Temperature in degrees Celsius
Patm	Surface atmospheric pressure in atm
P	Hydrostatic pressure in bar
pH	pH
CO2	CO2 concentration (mol/kg-soln)
pCO2	"standard" pCO2, CO2 partial pressure computed at in situ temperature and atmospheric pressure (μ atm)
fCO2	"standard" fCO2, CO2 fugacity computed at in situ temperature and atmospheric pressure (μ atm)
pCO2pot	"potential" pCO2, CO2 partial pressure computed at potential temperature and atmospheric pressure (μ atm)
fCO2pot	"potential" fCO2, CO2 fugacity computed at potential temperature and atmospheric pressure (μ atm)
pCO2insitu	"in situ" pCO2, CO2 partial pressure computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
fCO2insitu	"in situ" fCO2, CO2 fugacity computed at in situ temperature and total pressure (atm + hydrostatic) (μ atm)
HCO3	HCO3 concentration (mol/kg-soln)
CO3	CO3 concentration (mol/kg-soln)
DIC	DIC concentration (mol/kg-soln)
ALK	ALK, total alkalinity (mol/kg-soln)
OmegaAragonite	Omega aragonite, aragonite saturation state
OmegaCalcite	Omega calcite, calcite saturation state
NH4	NH4 concentration (mol/kg-soln)
NH3	NH3 concentration (mol/kg-soln)

B0H3	B(OH)3 concentration (mol/kg-soln)
B0H4	B(OH)4 concentration (mol/kg-soln)
H3P04	H3PO4 concentration (mol/kg-soln)
H2P04	H2PO4 concentration (mol/kg-soln)
HP04	HPO4 concentration (mol/kg-soln)
P04	PO4 concentration (mol/kg-soln)
H2S	H2S concentration (mol/kg-soln)
HS	HS concentration (mol/kg-soln)
SiOH4	SiOH4 concentration (mol/kg-soln)
Si00H3	SiOOH3 concentration (mol/kg-soln)
Si02OH2	SiO2OH2 concentration (mol/kg-soln)
HF	HF concentration (mol/kg-soln)
F	F concentration (mol/kg-soln)
HS04	HSO4 concentration (mol/kg-soln)
SO4	SO4 concentration (mol/kg-soln)
H	H concentration at specified pH scale (mol/kg-soln)
OH	OH concentration (mol/kg-soln)
NH4t	Supplied NHt concentration (mol/kg-soln); values are recycled if necessary
BOR	Calculated total borate concentration (mol/kg-soln)
Pt	Supplied Pt concentration (mol/kg-soln); values are recycled if necessary
HSt	Supplied HSt concentration (mol/kg-soln); values are recycled if necessary
Sit	Supplied Sit concentration (mol/kg-soln); values are recycled if necessary
FLUO	Calculated total fluoride concentration (mol/kg-soln)
ST	Calculated total sulfate concentration (mol/kg-soln)
SIR	substrate inhibitor ratio: [HCO3]/[H_free] (mol: μ mol)
H_free	H ion concentration on free scale (μ mol/kg)
H_sws	H ion concentration on seawater scale (μ mol/kg)
H_t	H ion concentration on total scale (μ mol/kg)
pH_free	pH calculated on the free scale
pH_sws	pH calculated on the seawater scale
pH_t	pH calculated on the total scale

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2014)

Author(s)

Kimberlee Baldry, Lennart Bach and Jean-Pierre Gattuso, <gattuso@obs-vlfr.fr>

References

Bach L. T., 2015 Reconsidering the role of carbonate ion concentration in calcification by marine organisms. *Biogeosciences* **12**, 4939-4951.

Dickson A. G. and Riley J. P., 1979 The estimation of acid dissociation constants in seawater media from potentiometric titrations with strong base. I. The ionic product of water. *Marine Chemistry* **7**, 89-99.

Dickson A. G., 1990 Standard potential of the reaction: $\text{AgCl}(\text{s}) + 1/2\text{H}_2(\text{g}) = \text{Ag}(\text{s}) + \text{HCl}(\text{aq})$, and the standard acidity constant of the ion HSO_4^- in synthetic sea water from 273.15 to 318.15 K. *Journal of Chemical Thermodynamics* **22**, 113-127.

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO_2 measurements. *PICES Special Publication* **3**, 1-191.

Khoo H. K., Ramette R. W., Culberson C. H. and Bates R. G., 1977 Determination of Hydrogen Ion Concentration in Seawater from 5 to 40°C: Standard Potentials at Salinities from 20 to 45. *Analytical Chemistry* **22**, vol49 29-34.

Lee K., Tae-Wook K., Byrne R.H., Millero F.J., Feely R.A. and Liu Y-M, 2010 The universal ratio of the boron to chlorinity for the North Pacific and North Atlantoc oceans. *Geochimica et Cosmochimica Acta* **74** 1801-1811.

Lueker T. J., Dickson A. G. and Keeling C. D., 2000 Ocean pCO_2 calculated from dissolved inorganic carbon, alkalinity, and equations for K_1 and K_2 : validation based on laboratory measurements of CO_2 in gas and seawater at equilibrium. *Marine Chemistry* **70** 105-119.

Millero F. J., 1995. Thermodynamics of the carbon dioxide system in the oceans. *Geochimica Cosmochimica Acta* **59**: 661-677.

Millero F. J., 2010. Carbonate constant for estuarine waters. *Marine and Freshwater Research* **61**: 139-142.

Millero F. J., Graham T. B., Huang F., Bustos-Serrano H. and Pierrot D., 2006. Dissociation constants of carbonic acid in seawater as a function of salinity and temperature. *Marine Chemistry* **100**, 80-84.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2014. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences Discussions* **11**, 5327-5397.

Perez F. F. and Fraga F., 1987 Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry* **21**, 161-168.

Roy R. N., Roy L. N., Vogel K. M., Porter-Moore C., Pearson T., Good C. E., Millero F. J. and Campbell D. M., 1993. The dissociation constants of carbonic acid in seawater at salinities 5 to 45 and temperatures 0 to 45°C. *Marine Chemistry* **44**, 249-267.

Schockman, K.M., Byrne, R.H., 2021. Spectrophotometric determination of the bicarbonate dissociation constant in seawater, *Geochimica et Cosmochimica Acta*.

Uppstrom L.R., 1974 The boron/chlorinity ratio of the deep-sea water from the Pacific Ocean. *Deep-Sea Research I* **21** 161-162.

Waters, J., Millero, F. J., and Woosley, R. J., 2014. Corrigendum to "The free proton concentration scale for seawater pH", [MARCHÉ: 149 (2013) 8-22], Mar. Chem. 165, 66-67.

Weiss, R. F., 1974. Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, **2**, 203-215.

Weiss, R. F. and Price, B. A., 1980. Nitrous oxide solubility in water and seawater, *Mar. Chem.*, **8**, 347-359.

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

Examples

```
## With a couple of variables
sir_full(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=0, Sit=0,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74", gas="potential", NH4t=0, HSt=0)

## With a couple of variables and non-zero nutrient concentrations
sir_full(flag=8, var1=8.2, var2=0.00234, S=35, T=25, P=0, Patm=1.0, Pt=5e-6, Sit=2e-6,
pHscale="T", kf="pf", k1k2="l", ks="d", b="u74", gas="potential", NH4t=10e-6, HSt=0.1e-6)

## Using vectors as arguments
flag <- c(8, 2, 8)
var1 <- c(8.2, 7.477544e-06, 8.2)
var2 <- c(0.002343955, 0.001649802, 2400e-6)
S <- c(35, 35, 30)
T <- c(25, 25, 30)
P <- c(0, 0, 0)
Pt <- c(0, 0, 0)
Sit <- c(0, 0, 0)
kf <- c("pf", "pf", "pf")
k1k2 <- c("l", "l", "l")
pHscale <- c("T", "T", "T")
b <- c("l10", "l10", "l10")
gas <- c("potential", "potential", "potential")
NH4t <- c(0, 0, 0)
HSt <- c(0, 0, 0)
sir_full(flag=flag, var1=var1, var2=var2, S=S, T=T, P=P, Pt=Pt, Sit=Sit,
kf=kf, k1k2=k1k2, pHscale=pHscale, b=b, gas=gas, NH4t=NH4t, HSt=HSt)

## Test with all flags
flag <- c((1:15), (21:25))
var1 <- c(8.200000, 7.308171e-06, 7.308171e-06, 7.308171e-06,
8.2, 8.2, 8.2, 8.2, 0.001646857, 0.001646857, 0.001646857, 0.0002822957,
0.0002822957, 0.00234, 258.2164, 258.2164, 258.2164, 258.2164, 258.2164 )
var2 <- c(7.308171e-06, 0.001646857, 0.0002822957, 0.00234, 0.001936461,
0.001646857, 0.0002822957, 0.00234, 0.001936461, 0.0002822957,
0.00234, 0.001936461, 0.00234, 0.001936461, 0.001936461, 8.2,
0.001646857, 0.0002822957, 0.00234, 0.001936461)
sir_full(flag=flag, var1=var1, var2=var2)
```

Description

Converts from practical to absolute salinity based on total alkalinity as well as on the concentrations of dissolved inorganic carbon, nitrate and silicate.

Usage

```
sp2sa_chem(SP, TA=2300e-6, DIC=2000e-6, NO3=0, SIOH4=0)
```

Arguments

SP	Practical salinity on the practical salinity scale
TA	Total alkalinity, in mol/kg, default is 2300 $\mu\text{mol/kg}$
DIC	Dissolved inorganic carbon concentration in mol/kg, default is 2000 $\mu\text{mol/kg}$
NO3	Total nitrate concentration in mol/kg, default is 0
SIOH4	Total silicate concentration in mol/kg, default is 0

Details

Converts from practical to absolute salinity from carbonate system parameters and ion concentration which mostly affect water density anomalies.

Value

SA	Absolute salinity (g/kg)
----	--------------------------

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., Wright D. G. and Millero F. J., 2011. The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* **7**, 363-387.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

sa2sp_chem does the reverse, sp2sa_geo

Examples

```
# Calculate the absolute salinity of a sample with practical Salinity of 35,
# Total Alkalinity of 0.00234 mol/kg and DIC of 0.00202 mol/kg
SA <- sp2sa_chem(SP=35, TA=0.00234, DIC=0.00202)
```

Description

Converts from practical to absolute salinity based on depth and geographic location.

Usage

```
sp2sa_geo(SP, P=0, long=1.e20, lat=1.e20)
```

Arguments

SP	Practical salinity on the practical salinity scale
P	Sea water pressure in dbar
long	Longitude in decimal degrees [0 ... +360] or [-180 ... +180]
lat	Latitude in decimal degrees [-90 ... 90]

Details

This function is almost an alias of function gsw_SA_from_SP of the gsw package on which it relies. The only difference is in that depth and location are optional. If location is not given, or incomplete (either longitude or latitude missing), an arbitrary location is chosen: the mid equatorial atlantic ocean. Note that this implies an error on computed SA ranging from 0 up to 0.02 g/kg.

Value

SA	Absolute salinity (g/kg)
----	--------------------------

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

sa2sp_geo does the reverse, sp2sa_chem

Examples

```
# Calculate the absolute salinity of a sample whose practical Salinity is 35,
# depth is 10 dbar and location is 188 degrees East and 4 degrees North.
SA <- sp2sa_geo(35, 10, 188, 4)      # 34.711778344814114
```

speciation

ionic forms as a function of pH

Description

Estimates the concentration of the various ionic forms of a molecule as a function of pH

Usage

```
speciation(K1=K1(), K2=NULL, K3=NULL, pH, conc=1)
```

Arguments

K1	First dissociation constant
K2	Second dissociation constant, default is NULL
K3	Third dissociation constant, default is NULL
pH	pH value, default is 8
conc	concentration of molecule in mol/kg, default is 1 mol/kg

Value

The function returns a data frame containing the following concentrations (in mol/kg if conc is given in mol/kg):

C1	ionic form 1, univalent, bivalent and trivalent molecules
C2	ionic form 2, univalent, bivalent and trivalent molecules
C3	ionic form 3, bivalent and trivalent molecules
C4	ionic form 4, trivalent molecules

Author(s)

Karline Soetaert <K.Soetaert@nioo.knaw.nl>

References

Zeebe R. E. and Wolf-Gladrow D. A., 2001 *CO₂ in seawater: equilibrium, kinetics, isotopes*. Amsterdam: Elsevier, 346 pp.

See Also

[bjerrum](#).

Examples

```

## Speciation of divalent species; example to estimate the various ionic forms
## of dissolved inorganic carbon (DIC = 0.0021 mol/kg) at a salinity of 35,
## a temperature of 25oC and an hydrostatic pressure of 0:
spec <- speciation (K1(35, 25, 0), K2(35, 25, 0), pH=8, conc=0.0021)
## where (spec\[C1]=[CO2], spec\[C2]=[HCO3-], spec\[C3]=[CO3--])

## Speciation of trivalent species (e.g., H3PO4, H2PO4-, HPO4--, PO4---)
speciation(K1p(), K2p(), K3p(), conc=0.001)

## Effect of temperature on pCO2 - Figure 1.4.18 of Zeebe and Wolf-Gladrow (2001)
Tseq <- seq(0, 30, by=0.5)
pHseq <- carb(flag=15, var1=2300e-6, var2=1900e-6, S=35, T=Tseq, P=0)$pH
CO2 <- speciation(K1(T=Tseq), K2(T=Tseq), conc=1900, pH=pHseq)$C1
pCO2 <- CO2/K0(T=Tseq)
plot(Tseq, pCO2, xlab="Temperature (oC)", ylab="pCO2 (uatm)", type="l",
main="effect of temperature on pCO2")
legend("topleft", c(expression(sum(CO[2]) == 1900~umol~kg^-1"),
expression(TA == 2300~umol~kg^-1)))

```

teos2eos_chem

Convert temperature and salinity from TEOS-10 to EOS-80

Description

Converts conservative temperature to in situ temperature and absolute salinity to practical salinity (SP). Salinity conversion depends on total alkalinity as well as on the concentrations of dissolved inorganic carbon, nitrate and silicate.

Usage

```
teos2eos_chem(SA, CT, P=0, TA=2300e-6, DIC=2000e-6, NO3=0, SiOH4=0)
```

Arguments

SA	Absolute salinity in g/kg
CT	Conservative temperature in degrees C
P	Sea water pressure in dbar
TA	Total alkalinity, in mol/kg, default is 2300 μ mol/kg
DIC	Dissolved inorganic carbon concentration in mol/kg, default is 2000 μ mol/kg
NO3	Total nitrate concentration in mol/kg, default is 0
SiOH4	Total silicate concentration in mol/kg, default is 0

Details

Conversion from absolute to practical salinity depends on carbonate system parameters and ion concentration which mostly affect water density anomalies.

Value

The function returns a data frame containing the following columns:

T	In situ temperature (deg C)
SP	Practical salinity (psu)

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., Wright D. G. and Millero F. J., 2011. The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* **7**, 363-387.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

eos2teos_chem, teos2eos_geo, sa2sp_cham, package gsw

Examples

```
# Calculate in situ temperature and practical salinity of a sample with
# Absolute salinity of 35 g/kg, Conservative temperature of 18 deg C,
# at 0 dbar and Total alkalinity of 0.00234 mol/kg and DIC of 0.00202 mol/kg
f <- teos2eos_chem(SA=35, CT=18, P=0, TA=0.00234, DIC=0.00202)
T <- f$T      # insitu temperature
SP <- f$SP     # Practical salinity
```

teos2eos_geo

Convert temperature and salinity from TEOS-10 to EOS-80

Description

Converts conservative temperature to in situ temperature and absolute salinity to practical salinity (SP). Salinity conversion depends on depth and geographic location.

Usage

```
teos2eos_geo(SA, CT, P=0, long=1.e20, lat=1.e20)
```

Arguments

SA	Absolute salinity in g/kg
CT	Conservative temperature in degrees C
P	Sea water pressure in dbar
long	Longitude in decimal degrees [0 ... +360] or [-180 ... +180]
lat	Latitude in decimal degrees [-90 ... 90]

Details

Conversion from absolute to practical salinity depends on water density anomaly which is correlated with silicate concentration. This function relies on silicate concentration taken from WOA (World Ocean Atlas) to evaluate the density anomaly.

Value

The function returns a data frame containing the following columns:

T	In situ temperature (deg C)
SP	Practical salinity (psu)

Author(s)

Jean-Marie Epitalon

References

McDougall T. J., Jackett D. R., Millero F. J., Pawlowicz R. and Barker P. M., 2012. A global algorithm for estimating Absolute Salinity. *Ocean Science* **8**, 1123-1134.

Pawlowicz R., Wright D. G. and Millero F. J., 2011. The effects of biogeochemical processes on oceanic conductivity/salinity/density relationships and the characterization of real seawater. *Ocean Science* **7**, 363-387.

Pawlowicz R., 2013. What every oceanographer needs to know about TEOS-10 (The TEOS-10 Primer). <http://www.teos-10.org/>

See Also

eos2teos_geo does the reverse, teos2eos_chem, sa2sp_geo, package gsw

Examples

```
# Calculate in situ temperature and practical salinity of a sample with
# Absolute salinity of 35 g/kg, conservative temperature of 18 deg C,
# depth is 10 dbar and location is 188 degrees East and 4 degrees North.
f <- teos2eos_geo(SA=35, CT=18, P=10, long=188, lat=4)
T <- f$T      # in situ temperature
SP <- f$SP     # Practical salinity
```

theta	<i>Potential temperature of seawater</i>
-------	--

Description

Computes theta, the potential temperature of seawater given original temperature, salinity, pressure, and reference pressure

Usage

```
theta(S=35, T=25, P=0, Pref=0)
```

Arguments

S	Salinity on the practical salinity scale, default is 35
T	Temperature in degrees Celsius, default is 25oC
P	Hydrostatic pressure in bar (surface = 0; 1000 db = 100 bar), default is 0
Pref	Reference hydrostatic pressure in bar, default is 0

Details

Computes the potential temperature of seawater relative to a chosen reference pressure following Fofonoff and Millard (1983). The potential temperature θ is the temperature that a water parcel would have if were moved adiabatically to another pressure level $Pref$. Typically, the potential temperature is referenced to the surface ($Pref = 0$). The potential teperature depends on the original salinity S , *in-situ* temperature T and pressure P .

This routine is essentially a wrapper for the [swTheta](#) routine of the 'oce' package. Unlike the latter, pressure units here are given in bars and method="unesco" is prescribed.

Value

theta	potential temperature of seawater (C)
-------	---------------------------------------

Author(s)

James Orr <james.orr@lsce.ipsl.fr>

References

Fofonoff, P. and R. C. Millard Jr, 1983. Algorithms for computation of fundamental properties of seawater. *Unesco Technical Papers in Marine Science*, **44**, 53 pp.

See Also

[swTheta](#).

Examples

```
#Calculate the potential temperature for a sample at 1000 db referenced to the surface
theta <- theta(S=35, T=25, P=100, Pref=0)
```

tris

pH of TRIS buffer

Description

Calculates the pH value of TRIS buffered artificial seawater solutions (on the total scale in mol/kg-soln)

Usage

```
tris(S=35,T=25,b=0.04,k="d98",warn="y")
```

Arguments

S	Salinity, default is 35
T	Temperature in degrees Celsius, default is 25oC
b	Molality of TRIS/TRISH+ in moles per kg of water, default is 0.04 mol/kg-H2O
k	"d98" for DelValls and Dickson 1998, "m18" for using tris characterization by Mueller et al (2018), default is "d98"
warn	"y" to show warnings when S,T and/or b go beyond the valid range for the chosen k; "n" to suppress warnings. The default is "y".

Details

The models used to calculate the return value of this function are based on experimental data. It is critical to consider that each formulation refers to the artificial seawater solution applied during the characterization experiment and is only valid for the studied ranges of temperature and salinity:

- DelValls and Dickson (1998): S ranging between 20 and 40, T ranging between 0 and 45oC, and b being 0.04 mol/kg-H2O.
- Mueller et al. (2018): S ranging between 5 and 40, T ranging between 5 and 45oC, and b ranging between 0.01 and 0.04 mol/kg-H2O.

Note that the arguments can be given as a unique number or as vectors. If the lengths of the vectors are different, the longer vector is retained and only the first value of the other vectors is used. It can therefore be critical to use vectors of the same length.

Value

tris The function returns the pH value of TRIS buffered artificial seawater solutions (on the total scale in mol/kg-soln)

Author(s)

Jean-Pierre Gattuso <gattuso@obs-vlfr.fr> and Jens Daniel Mueller <jens.mueller@io-warnemuende.de>

References

DelValls, T. A., and Dickson, A. G., 1998 The pH of buffers based on 2-amino-2-hydroxymethyl-1,3-propanediol ('tris') in synthetic sea water. *Deep Sea Research Part I: Oceanographic Research Papers* **45(9)**, 1541-1554. [https://doi.org/10.1016/S0967-0637\(98\)00019-3](https://doi.org/10.1016/S0967-0637(98)00019-3)

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO₂ measurements. *PICES Special Publication 3*, 1-191.

Mueller, J. D., Bastkowski, F., Sander, B., Seitz, S., Turner, D. R., Dickson, A. G., and Rehder, G., 2018 Metrology for pH measurements in brackish waters- Part 1: Extending electrochemical pHT measurements of TRIS buffers to salinities 5-20. *Frontiers in Marine Science* **5:176**, 1-12. <https://doi.org/10.3389/fmars.2018.00176>

See Also

[amp](#), [pHslope](#), [pH](#).

Examples

```
##Example from Mueller et al. (2018), should give test value pHT = 8.0703
tris(S=20,T=25,b=0.04,k="m18")
```

vapress

Computes vapor pressure of seawater

Description

Computes vapor pressure of seawater (atm) from temperature and salinity

Usage

```
vapress(S=35, T=25, form="d2007")
```

Arguments

S Salinity on the practical salinity scale, default is 35

T Temperature in degrees Celsius, default is 25oC

form choose either "d2007" for the best-practices formulation to compute vapor pressure of seawater from Dickson et al. (2007) or "wp1980" for the formulation from Weiss and Price (1980).

Details

Computes the vapor pressure of seawater pH20 following best practices (Dickson et al., 2007). That computed pH20 is identical, when rounded to the 4th decimal place, with that computed by the equation from Weiss and Price (1980).

Value

vapress Vapor pressure of seawater in atm

Author(s)

James Orr <james.orr@lsce.ipsl.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R. (2007) Guide to best practices for ocean CO₂ measurements. *PICES Special Publication* **3**, 1-191.

Weiss, R. F. (1974) Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, **2**, 203-215.

Weiss, R. F. and Price, B. A. (1980) Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

See Also

[x2pCO2](#), and [p2xC02](#).

Examples

```
pH20 <- vapress(S=35, T=25, form="d2007")
```

x2pCO2

Converts mole fraction to partial pressure of CO₂

Description

Converts xCO₂ (mole fraction of CO₂) into pCO₂ (partial pressure of CO₂)

Usage

```
x2pCO2(S=35, T=25, Patm=1.0, xCO2=400)
```

Arguments

S	Salinity on the practical salinity scale, default is 35
T	Temperature in degrees Celsius, default is 25oC
Patm	Atmospheric pressure in atmospheres, default is 1.0
xCO2	Mole fraction of CO ₂ in ppm, default is 400

Details

The mole fraction xCO₂ (ppm) is computed from pCO₂ (μ atm) using the following equation: $xCO_2 = pCO_2(Patm - pH2O)$, where pH2O is the vapor pressure of seawater computed following best practices (Dickson et al., 2007). That computed pH2O is identical, when rounded to the 4th decimal place, with that computed by the equation from Weiss and Price (1980).

Value

pCO2 Partial pressure of CO2 in μatm .

Note

Warning: pCO2 estimates below 100 m are subject to considerable uncertainty. See Weiss (1974) and Orr et al. (2015)

Author(s)

James Orr <james.orr@lsce.ipsl.fr>

References

Dickson A. G., Sabine C. L. and Christian J. R., 2007 Guide to best practices for ocean CO2 measurements. *PICES Special Publication* **3**, 1-191.

Orr J. C., Epitalon J.-M. and Gattuso J.-P., 2015. Comparison of seven packages that compute ocean carbonate chemistry. *Biogeosciences* **12**, 1483-1510.

Weiss, R. F. (1974) Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Marine Chemistry*, **2**, 203-215.

Weiss, R. F. and Price, B. A. (1980) Nitrous oxide solubility in water and seawater, *Marine Chemistry*, **8**, 347-359.

See Also

[p2xCO2](#) and [vapress](#)

Examples

```
## Atmospheric pressure is rarely equal to 1 atm exactly
## Over the Southern Ocean Patm=0.97 is more realistic
pCO2_socn <- x2pCO2(S=35, T=0, Patm=0.97, xCO2=400.0)
print(pCO2_socn)
## The result (385.6322 uatm) is 12 uatm less than if it was wrongly assumed that Patm=1.0

## Show effect of temperature on pCO2 computed from xCO2, and on resulting variables from "carb"
S <- 35
ALK <- 2300e-6
T <- seq(0,30,5)
xCO2 <- 400
pCO2 <- x2pCO2(S=35, T=T, Patm=1, xCO2=400)
results <- carb(flag=24, var1=pCO2, var2=ALK, S=S, T=T, P=0, Pt=0, Sit=0,
  pHscale="T", kf="pf", k1k2="1", ks="d", b="u74")
print(results)
```

Index

- * **datasets**
 - alkalinity, 3
 - kfg, 75
 - Pcoeffs, 102
 - seacarb_test_P0, 138
 - seacarb_test_P300, 139
 - seaFET, 139
- * **utilities**
 - amp, 4
 - bjerrum, 5
 - bor, 7
 - buffer, 8
 - buffergen, 13
 - buffesm, 18
 - carb, 23
 - carbb, 29
 - carbfull, 35
 - d2p, 41
 - derivnum, 42
 - eos2teos_chem, 47
 - eos2teos_geo, 48
 - errors, 50
 - f2pCO2, 57
 - fCO2insi, 58
 - K0, 59
 - K1, 60
 - K1p, 63
 - K2, 64
 - K2p, 67
 - K2si, 68
 - K3p, 69
 - Kb, 71
 - kconv, 72
 - Kf, 74
 - Khs, 76
 - Kn, 77
 - Ks, 78
 - Ksi, 79
 - Kspa, 80
- Kspc, 82
- Kw, 83
- oa, 84
- Om, 89
- p2d, 93
- p2fCO2, 94
- p2xCO2, 95
- pCa, 96
- pCO2insi, 101
- Pcorrect, 103
- pgas, 104
- pH, 109
- pHconv, 110
- pHinsi, 111
- pHslope, 115
- pHspec, 116
- pmix, 117
- ppH, 121
- psi, 126
- pTA, 130
- rho, 135
- sa2sp_chem, 135
- sa2sp_geo, 137
- sf_calc, 140
- sf_calib, 141
- sir, 143
- sir_b, 149
- sir_full, 155
- sp2sa_chem, 161
- sp2sa_geo, 163
- speciation, 164
- teos2eos_chem, 165
- teos2eos_geo, 166
- theta, 168
- tris, 169
- vapress, 170
- x2pCO2, 171

- alkalinity, 3
- amp, 4, 109, 115, 117, 170

bjerrum, 5, 164
 bor, 7
 buffer, 8, 126
 buffergen, 13
 buffesm, 18
 carb, 23, 89
 carbb, 29
 carbfull, 35
 d2p, 41, 93
 derivnum, 42
 eos2teos_chem, 47
 eos2teos_geo, 48
 errors, 50
 f2pCO2, 57, 59, 94, 101
 fCO2insi, 58, 101
 K0, 59
 K1, 60, 66
 K1p, 63, 68, 70
 K2, 63, 64
 K2p, 64, 67, 70
 K2si, 68
 K3p, 64, 68, 69
 Kb, 71
 kconv, 72, 111
 Kf, 74
 kfg, 75
 Khs, 76
 Kn, 77
 Ks, 78
 Ks_i, 79
 Kspa, 80, 83
 Kspc, 81, 82
 Kw, 83
 matplot, 6
 oa, 84
 Om, 89
 p2d, 42, 93
 p2fCO2, 58, 94
 p2xCO2, 95, 171, 172
 par, 5, 6
 pCa, 96
 pCO2insi, 59, 101
 Pcoeffs, 102
 Pcorrect, 103
 pgas, 89, 104
 pH, 4, 109, 115, 117, 170
 pHconv, 73, 110
 pHinsi, 111
 pHslope, 4, 109, 115, 117, 170
 pHspec, 116
 plot.default, 5
 pmix, 89, 117
 ppH, 89, 121
 psi, 126
 pTA, 89, 130
 rho, 135
 sa2sp_chem, 135
 sa2sp_geo, 137
 seacarb_test_P0, 138
 seacarb_test_P300, 139
 seaFET, 139
 sf_calc, 140, 142
 sf_calib, 141, 141
 sir, 143
 sir_b, 149
 sir_full, 155
 sp2sa_chem, 161
 sp2sa_geo, 163
 speciation, 6, 130, 164
 swTheta, 168
 teos2eos_chem, 165
 teos2eos_geo, 166
 theta, 168
 title, 5
 tris, 4, 109, 115, 117, 169
 vapress, 96, 170, 172
 x2pCO2, 96, 171, 171